



# Recent results on the dark sector from Belle II


Paul Feichtinger, on behalf of the Belle II Collaboration

**PASCOS** - 27th International Symposium on Particles, Strings and Cosmology

Heidelberg, 26.07.2022


# Introduction

- B-factories have unique reach in direkt searches for the light dark sector
  - low mass mediator particles on the MeV-GeV scale
- Recent results from Belle II:
  - Dark Higgsstrahlung
  - $Z' \rightarrow \text{invisible}$
  - $\text{TT}$  resonance
    - $Z' \rightarrow \text{TT}$
    - $S \rightarrow \text{TT}$
    - $\text{ALP} \rightarrow \text{TT}$




# Belle II and SuperKEKB

- B-factory located in Tsukuba, Japan
- colliding electrons and positrons at  $m_{Y(4S)} = 10.58 \text{ GeV}/c^2$
- collected luminosity from 2019-2022:  $424 \text{ fb}^{-1}$
- peak luminosity world record:  $4.7 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
- target x50 Belle data ( $\approx 50 \text{ ab}^{-1}$ )



SuperKEKB accelerator




# Belle II and SuperKEKB

see talk by Doris Kim on  
Wednesday (session B)


- general purpose detector: B and D physics, quarkonium,  $\tau$ -physics, dark sector, ...
- clean collision environment ( $e^+ \leftrightarrow e^-$ )
- large solid angle coverage ( $> 90\%$ )
  - well known missing mass and energy
- excellent PID
- dedicated low-multiplicity triggers
  - two/three-track trigger
  - $E_{\text{ECL}} > 1 \text{ GeV}$  trigger
  - single muon trigger (drift chamber + muon detector)

Belle II detector



# Dark Higgsstrahlung: $e^+e^- \rightarrow A' h'$


- U(1)' extension to SM » [Phys. Rev. D 79, 115008 \(2009\)](#)
  - Dark photon  $A'$ 
    - coupled to SM photon via kinetic mixing parameter  $\epsilon$
    - mass generated via spontaneous symmetry breaking
  - Dark Higgs  $h'$ 
    - couples with  $\alpha_D$  to  $A'$
    - does not mix with SM Higgs



- $M_{h'} < M_{A'}$ :  $h'$  is long-lived (invisible)  $\Rightarrow$  2 charged tracks

$M_{A'} \Leftrightarrow M_{\mu\mu}$  and  $M_{h'} \Leftrightarrow M_{\text{recoil}}$

partially constrained by [KLOE \(2015\)](#)





# Dark Higgsstrahlung

- data
  - 8.34 fb<sup>-1</sup> (2019)
- backgrounds
 

○ e <sup>+</sup> e <sup>-</sup> → μ <sup>+</sup> μ <sup>-</sup> (γ)	79%
○ e <sup>+</sup> e <sup>-</sup> → τ <sup>+</sup> τ <sup>-</sup> (γ)	18%
○ e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> μ <sup>+</sup> μ <sup>-</sup>	3%
- selection
  - two reconstructed muons, p<sub>T</sub><sup>μμ</sup> > 0.1 GeV/c
  - recoil momentum in the ECL barrel, no nearby photon
  - cut on helicity angle
- strategy
  - scan for excess in 2D plane of M<sub>recoil</sub> vs M<sub>μμ</sub>
  - ~9000 rotated elliptical mass windows to test signal hypotheses

→ observed yields →




# Dark Higgsstrahlung - Results



submitted to PRL

» [ArXiv: 2207.00509](https://arxiv.org/abs/2207.00509)


- no significant excess above background was observed → 90% CL upper limits



world leading limits for  $1.65 < M_{A'} < 10.51 \text{ GeV}/c^2$



# Z': the $L_\mu - L_\tau$ model



- extension of standard model with a  $U(1)'$  group
- gauging  $L_\mu - L_\tau$ , the difference of leptonic  $\mu$  and  $\tau$  number
- $Z'$  is resulting new massive gauge boson that couples only to  $\mu$  and  $\tau$  leptons
- can provide solution for
  - dark matter puzzle ( $Z'$  as mediator between SM and DS)
  - $(g-2)_\mu$
  - $b \rightarrow s\mu\mu, R_K, R_{K^*}$  anomalies

$$\mathcal{L} = \sum_{\ell} \theta g' \bar{\ell}' \gamma^\mu Z'_\mu \ell$$

» [Altmannshofer et al. JHEP 1612 \(2016\) 106](#)  
 » [Shuve et al. PRD 89, 113004 \(2014\)](#)

- Belle II search for  $Z'$  in  $\mu^+\mu^-$  final state with



- $Z' \rightarrow$  invisible (neutrinos / dark matter)
- $Z' \rightarrow \tau\tau$

final states with missing energy,  $M_{Z'} \leftrightarrow M_{\text{recoil}}$





# Z' → invisible

- data
  - 79.7 fb<sup>-1</sup> (2019-2020)
- backgrounds
  - e<sup>+</sup>e<sup>-</sup> → μ<sup>+</sup>μ<sup>-</sup>(γ)
  - e<sup>+</sup>e<sup>-</sup> → τ<sup>+</sup>τ<sup>-</sup>(γ)
  - e<sup>+</sup>e<sup>-</sup> → e<sup>+</sup>e<sup>-</sup>μ<sup>+</sup>μ<sup>-</sup>
- selection
  - two reconstructed muons, p<sub>T</sub><sup>μμ</sup> > 0.4 GeV/c
  - recoil momentum in the ECL barrel, no nearby photon
  - neural network trained to optimize Punzi FOM
  - » [Eur. Phys. J. C 82, 121 \(2022\)](#)
- strategy
  - template fit in 2D plane of θ<sub>recoil</sub><sup>CMS</sup> vs M<sub>recoil</sub><sup>2</sup>



# $Z'$ $\rightarrow$ invisible - Results



- no significant excess above background was observed  $\rightarrow$  90% CL upper limits



excluded fully invisible  $Z'$  as explanation for  $(g-2)_\mu$  for  $0.8 < M_{Z'} < 5.0 \text{ GeV}/c^2$




# Z', S, ALP $\rightarrow$ $\tau\tau$

- search for a  $\tau\tau$  resonance in  $\mu^+\mu^-\tau^+\tau^-$  final states
- data
  - $63.3 \text{ fb}^{-1}$  (2019-2020)
- selection
  - 4 tracks:  $2\mu + 2 e/\mu/\pi$  (1-prong  $\tau$  decay)
  - $M(4\text{-track}) < 9.5 \text{ GeV}/c^2$
  - 8 neural networks - trained for different ranges in  $M_{\text{recoil}}(\mu\mu)$
- backgrounds
  - $e^+e^- \rightarrow \tau^+\tau^-(\gamma)$  (1x3 prong)
  - $e^+e^- \rightarrow qq$  ( $q=u,d,s,c$ )
  - $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$
  - $e^+e^- \rightarrow \mu^+\mu^-\tau^+\tau^-$  ← no ISR in simulation
  - $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$
  - $e^+e^- \rightarrow \mu^+\mu^-\pi^+\pi^-, e^+e^- \rightarrow e^+e^-\chi_{\text{hadronic}}$  ← not simulated
- strategy
  - fit for a signal in  $M_{\text{recoil}}$  above floating background



# Z', S, ALP $\rightarrow$ $\tau\tau$ - Results

- no significant excess above background was observed  $\rightarrow$  90% CL upper limits




first constraints on S for  $M_S > 6.5 \text{ GeV}/c^2$  + first direct constraints for ALP  $\rightarrow$   $\tau\tau$

- $\gg$  [1] [PRD 95 \(2017\) 075003](#)
- $\gg$  [2] [arXiv:2110.10698](#)

# Summary

- Belle II recorded  $424 \text{ fb}^{-1}$  so far → only partially used for present results
- new results for
  - Dark Higgsstrahlung search
  - $Z' \rightarrow$  invisible search
  - $Z', S, \text{ALP} \rightarrow \pi\pi$  search
- suitable for light dark sector searches
  - hermetic detector
  - clean collision environment
  - excelled particle identification
  - dedicated low multiplicity triggers
- more to come in the future



Thank you!

backup slides ↴




# From KEKB to SuperKEKB

## Belle + KEKB (1999-2010)

- peak luminosity:  $2.1 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
- collected almost  $1 \text{ ab}^{-1}$  at different resonances and off-resonances

## Belle II + SuperKEKB (first collisions in 2019)


- **nanobeam scheme** + **increased beam current**  
→ goal is 30 times higher luminosity
- luminosity world record ( $2.9 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ )
- goal: collect  $50 \text{ ab}^{-1}$  during lifetime (now:  $213 \text{ fb}^{-1}$ )
- challenges: dealing with higher machine backgrounds and trigger rates





challenge: increased beam backgrounds

(LER/HER)	E (GeV)	$\beta_y^*$ (mm)	$\beta_x^*$ (cm)	$\varphi$ (mrad)	I (A)	L ( $\text{cm}^{-2} \text{ s}^{-1}$ )
KEKB	3.5/8.0	5.9/5.9	120/120	11	1.6/1.2	$2.1 \times 10^{34}$
SuperKEKB	4.0/7.0	0.27/0.30	3.2/2.5	41.5	3.6/2.6	$60 \times 10^{34}$

# Trigger




+ neural triggers (single track)






# Dark Higgsstrahlung

- $M_{h'} > M_{A'}$ :  $h' \rightarrow A'A' \Rightarrow$  6 charged tracks searches by [BaBar \(2012\)](#) and [Belle \(2015\)](#)
- 2-track trigger
- control samples
  - $\mu\mu\gamma$   $\mu\mu(\gamma)$  background
  - $e\mu$   $\tau\tau$  background





systematics





source	uncertainty	target
Pre-selections	<b>2 - 9.1%</b>	BKG & signal
BKG shape	<b>9.3%</b> (region specific)	BKG
$C_\eta$ cut	<b>1%</b>	BKG
Mass resolution	<b>2.4%</b> (on average)	signal
Eff. Inside windows	<b>2 - 5%</b>	signal
Theory (BR $A'$ )	<b>4%</b>	signal


# Dark Higgsstrahlung






# Dark Higgsstrahlung



# $Z'$ → invisible




- within the  $L_\mu$ - $L_\tau$  model the  $Z'$  can decay invisibly only via neutrinos
- if we allow a hypothetical decay of the  $Z'$  to dark matter, the  $\text{BF}(Z' \rightarrow \text{invisible})$  can be enhanced
- we consider both cases in our search







# $Z'$ $\rightarrow$ invisible

- previous searches for  $Z' \rightarrow \mu^+\mu^-$  by [BaBar](#), [Belle](#), [CMS](#)
- 2-track trigger
- control samples
  - $\mu\mu\gamma$  selection+NN studies (low mass)
  - $e\mu$  selection+NN studies (medium + high mass)
  - $ee(\gamma)$   $\gamma$  veto studies





## systematics

Source	Low mass	Medium mass	High mass
selections	2.7%	6.5%	8.3%
Mass resolution	10%	10%	10%
Background shapes	3.2%	8.6%	25%
Photon veto	34%	5%	5%
luminosity	1%	1%	1%




# $Z', S, \text{ALP} \rightarrow \tau\tau$

- $L_\mu - L_\tau$ :  $Z'$ 
  - first search in  $\tau\tau$  final state
- leptophilic scalar:  $S$ 
  - partially constraint by BaBar in  $S \rightarrow \mu\mu$
  - first search in  $\tau\tau$  final state
- ALP:  $a$ 
  - assume  $C_{ee} = C_{\mu\mu} = C_{\tau\tau}$  and  $C_{\gamma\gamma} = C_{Z\gamma} = 0$
  - ALP- $\tau$  coupling unconstrained



# Z', S, ALP $\rightarrow$ $\tau\tau$

- control sample
  - $2\pi + 2e/\mu/\pi$
- 3-track or single muon trigger



$e^+e^- \rightarrow e^+e^- X_{\text{hadronic}}$

Missing ISR


## systematics

source	Uncertainty (%)
trigger	2.7
Particle ID	3.9-6.2
Tracking	3.6
Fit bias	4
MLP selection	2.8
Mass resolution	3
Efficiency interpolation	2.5
Luminosity	1
other	1
<b>Total</b>	<b>8.8-9.9</b>




# Punzi-Net

» [Eur. Phys. J. C 82, 121 \(2022\)](#)



The constants a and b are the number of sigmas corresponding to one-sided Gaussian tests at some predefined significance level,  $\alpha$  and  $\beta$ . Here  $\alpha$  is the probability of rejecting  $H_0$  when it is true (type I error), and  $\beta$  is the probability of not rejecting  $H_0$  when instead  $H_{s,0}$  is true (type II error).



**Trained on 4 kinematic variables:**


variable	description
$p_{i,\text{thrust}}^*(\mu)$	The transverse momentum component of the muons with respect to the thrust axis.
$p_{i,\mu_{\min}}^*(\mu_{\max})$	The transverse momentum component of the higher energetic muon with respect to the lower energetic muon.
$p_{i,\mu_{\min}}^*(\mu_{\max})$	The longitudinal momentum component of the higher energetic muon with respect to the lower energetic muon.
$p_i^*(\mu^+\mu^-)$	The transverse momentum of the dimuon system.

» [G. Punzi, Sensitivity of searches for new signals and its optimization](#)

# Projection of integrated luminosity delivered by SuperKEKB to Belle II

Target scenario: extrapolation from 2021 run including expected improvements.

Base scenario: conservative extrapolation of SuperKEKB parameters from 2021 run



- We start long shutdown I (LS1) from summer 2022 for 15 months to replace VXD. There will be other maintenance/improvement works of machine and detector.
- We resume physics running from Fall 2023.
- A SuperKEKB International Taskforce (aiming to conclude in summer 2022) is discussing additional improvements.
- An LS2 for machine improvements could happen on the time frame of 2026-2027