


Dark Sector searches at Belle II



Giacomo De Pietro

INFN Roma Tre




55th Rencontres de Moriond Electroweak Interactions and Unified Theories

20-27 March 2021 @ Home

SuperKEKB: a new Intensity Frontier machine

SuperKEKB is an upgraded B-factory located at KEK (Tsukuba, Japan)


It's an asymmetric e^+e^- collider operating mainly at **10.58 GeV** ($\Upsilon(4S)$, but foreseen runs from $\Upsilon(2S)$ to $\Upsilon(6S)$)



I (A): $\sim 1.6/1.2$


β_y^* (mm): $\sim 5.9/5.9$

nano-beam
scheme



I (A): $\sim 2.8/2.0$

β_y^* (mm): $\sim 0.27/0.3$



30x peak luminosity:
 $6 \cdot 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$

SuperKEKB: a new Intensity Frontier machine

SuperKEKB


located at

It's an asy

operating

($\Upsilon(4S)$), but fore



KEKB



$I(A): \sim 1.6/1.2$

$\beta_y^* (mm): \sim 5.9/5.9$

$\beta_y^* (mm): \sim 0.27/0.3$



luminosity:
6 · 10³⁵ cm⁻² s⁻¹

Belle II: a new Intensity Frontier detector

Electromagnetic Calorimeter (ECL):

CsI(Tl) crystals, waveform sampling to measure time, energy, and pulse-shape.

K_L and muon detector (KLM):

Resistive Plate Counters (RPC) (outer barrel)
Scintillator + WLSF + MPPC (endcaps, inner barrel)

Magnet:

1.5 T superconducting

Vertex detectors (VXD):

2 layer DEPFET pixel detectors (PXD)
4 layer double-sided silicon strip detectors (SVD)

Trigger rate limits:


Hardware: < 30 kHz
Software: < 10 kHz

Central drift chamber (CDC):

He(50%):C₂H₆ (50%), small cells,
fast electronics

Particle Identification (PID):

Time-Of-Propagation counter (TOP) (barrel)
Aerogel Ring-Imaging Cherenkov Counter (ARICH) (FWD)



SuperKEKB and Belle II operations


First collisions: 26th April 2018



Collected 0.5 fb^{-1} in 2018 (pilot run)

Collected more than
 100 fb^{-1} since 2019

Goal: collect up to 50 ab^{-1} !



Dark Matter


It is “dark”.

It exists...


A lot of experimental techniques
to probe the Dark Matter existence:

- production at colliders;
- production in fixed target experiments;
- indirect astronomic searches;
- direct underground searches.

Rotational curves of the galaxies



Bullet clusters




Dark Matter

It is “dark”.


It exists...

A lot of experimental techniques to probe the Dark Matter existence:

- production at colliders.



Dark Matter coupling to SM



Different possible portals between **Dark Matter** and **Standard Model** depending on the **dark mediator X**:


Vector portal → Dark Photon / Z'

Scalar portal → Dark Higgs / Dark Scalar

Pseudoscalar portal → Axion-Like Particles

Neutrino portal → Sterile Neutrinos

Dark Matter coupling to SM



Different possible portals between **Dark Matter** and **Standard Model** depending on the **dark mediator X**:

Vector portal \rightarrow Dark Photon / **Z'**

Scalar portal \rightarrow **Dark Higgs** / Dark Scalar

Pseudoscalar portal \rightarrow **Axion-Like Particles**

Neutrino portal \rightarrow Sterile Neutrinos

$L_\mu - L_\tau$ model: Z' to invisible


New light gauge boson Z' coupling only to 2nd and 3rd generation of leptons ($L_\mu - L_\tau$ model):

$$\mathcal{L} = \sum_{\ell = \mu, \tau, \nu_{\mu,L}, \nu_{\tau,L}} \theta g' \bar{\ell} \gamma^\mu Z'_\mu \ell \quad \text{Shuve et al. (2014), arXiv:1403.2727}$$

g' is the coupling constant introduced by the model

It may explain:

- DM puzzle;
- $(g-2)_\mu$ anomaly;
- $R(K)$ and $R(K^*)$ anomalies.



Invisible decay studied here for the first time

Looking for an invisibly decaying Z' produced with a pair of muons:

- Z' can decay to neutrinos or light DM if kinematically accessible;
- $\text{BF}(Z' \rightarrow \text{invisible}) = 1$ if $m_{Z'} < 2m_\mu$.


First physics paper by Belle II:
PRL 124 (2020) 141801

$L_\mu - L_\tau$ model: Z' to invisible

Measurement done using **2018 pilot run data**: only 276 pb^{-1} usable due to trigger conditions.

Looking for:

- a peak in the mass distribution of the system recoiling against the dimuon pair;
- nothing else in the rest of the event.




$L_\mu - L_\tau$ model: Z' to invisible


Measurement done using **2018 pilot run data**: only 276 pb^{-1} usable due to trigger conditions.

Looking for:

- a peak in the mass distribution of the system recoiling against the dimuon pair;
- nothing else in the rest of the event.





PRL 124 (2020) 141801



$L_\mu - L_\tau$ model: Z' to invisible

Short term projections including several improvements:

- much higher integrated luminosity (already on tape);
- analysis improvements (better muonID, MVA selection);
- new triggers w.r.t. 2018 pilot run.



Starting to probe the $(g-2)_\mu$ band already with 50 fb^{-1}


Axion-Like Particles: $a \rightarrow \gamma\gamma$

Axion-Like Particles (ALPs) are pseudoscalar particles (a) that couple to bosons.


Unlike QCD Axions, ALPs have no relation between mass and coupling.

Belle II focused on the **coupling to photons**:

$$\mathcal{L} \supset -\frac{g_{a\gamma\gamma}}{4} a F_{\mu\nu} \tilde{F}^{\mu\nu} \quad \tau_a \sim 1/g_{a\gamma\gamma}^2 m_a^3$$



Dolan et al. (2017), *arXiv:1709.00009*



Investigating the photon coupling $g_{a\gamma\gamma}$ in ALP-strahlung


First search at B-factories

Axion-Like Particles: $a \rightarrow \gamma\gamma$



Select fully neutral events consisting of 3 isolated photons with a total invariant mass consistent with $s^{1/2}$.

Search strategy optimized to maximize ALP sensitivity:

- low ALP mass \rightarrow **diphoton mass spectrum**; \dashrightarrow
- high ALP mass \rightarrow **recoil mass spectrum**.




PRL 125 (2020) 161806



Axion-Like Particles: $a \rightarrow \gamma\gamma$

Search conducted with 445 pb^{-1} of **2018 pilot run data**:

- 500 fits in sliding ranges with steps of half mass resolution;
- no excess observed (largest local significance: 2.8σ).





Axion-Like Particles: $a \rightarrow \gamma\gamma$

Search conducted with 445 pb⁻¹ of **2018 pilot run data**:

- 500 fits in sliding ranges with steps of half mass resolution;
- no excess observed (largest local significance: 2.8 σ).


$$\sigma_a = \frac{g_{a\gamma\gamma}^2 \alpha_{\text{QED}}}{24} \left(1 - \frac{m_a^2}{s}\right)^3$$



Dark Higgsstrahlung

Dark Photon (A') mass could be generated via a spontaneous symmetry breaking mechanism, adding a Dark Higgs boson (h') to the model:



- Dark Higgsstrahlung process: $e^+ e^- \rightarrow A'^* \rightarrow A' h'$



Dark Higgsstrahlung

Dark Photon (A') mass could be generated via a spontaneous symmetry breaking mechanism, adding a Dark Higgs boson (h') to the model:

- Dark Higgsstrahlung process: $e^+ e^- \rightarrow A'^* \rightarrow A' h'$



At Belle II we are exploring the invisible h' decay ($m_{h'} < m_{A'}$), constrained only by **KLOE**:

Babusci et al. (2015), arXiv:1501.06795

$$e^+ e^- \rightarrow A'^* \rightarrow A' (\rightarrow \mu^+ \mu^-) h' (\rightarrow \text{inv.})$$

Very promising results even considering only the 2019 data set (9 fb^{-1}):

- accessing an unconstrained region beyond the KLOE coverage;
- probing non-trivial $\epsilon^2 \alpha_D$ couplings.

Other ongoing Dark Sector searches @ Belle II

Dark Photon (A') decays

invisible decay *coming soon*

leptonic decays

Dark Scalar:

$$e^+ e^- \rightarrow \tau^+ \tau^- S (\rightarrow l^+ l^-)$$

Dark Scalar with displaced vertex:

$$B^+ \rightarrow K^+ S (\rightarrow l^+ l^-)$$

Inelastic Dark Matter

Other Z' decays:

$$e^+ e^- \rightarrow \mu^+ \mu^- Z' (\rightarrow \mu^+ \mu^-)$$

$$e^+ e^- \rightarrow \mu^+ \mu^- Z' (\rightarrow \tau^+ \tau^-)$$

Magnetic monopoles

... and many others!



Summary

- ✓ Belle II will lead the field in the Dark Sector searches in the coming years;
- ✓ We successfully started operations in 2018; up to now we collected more than 100 fb^{-1} of collisions data;
- ✓ First results with early data (2018 pilot run) are published:
 - Z' \rightarrow invisible search ([PRL 124 \(2020\) 141801](#));
 - ALP $\rightarrow \gamma\gamma$ search ([PRL 125 \(2020\) 161806](#));
- ✓ Next results in the pipeline using the data collected since 2019:
 - Dark Higgsstrahlung;
 - A' \rightarrow invisible search.



Thank you
for your
attention




Backup
slides

Taking data during the 2020 pandemic...

Non-stop operations with COVID-19 pandemic:

- social distancing requirements;
 - strong developments for close to or fully **remote sub-system operations**;
 - **huge commitments from japanese colleagues and residents in Japan**;
- only 40 persons on site from March to July.



Belle II
data taking
efficiency:
85%

courtesy of K. Matsuoka

SuperKEKB machine parameters

Parameter	KEKB Design	KEKB Achieved	SuperKEKB Design
Energy (GeV) (LER/HER)	3.5/8.0	3.5/8.0	4.0/7.0
β_y^* (mm)	10/10	5.9/5.9	0.27/0.30
β_x^* (mm)	330/330	1200/1200	32/25
ϵ_x (nm)	18/18	18/24	3.2/5.3
$\frac{\epsilon_y}{\epsilon_x}$ (%)	1	0.85/0.64	0.27/0.24
σ_y (μm)	1.9	0.94 $\xrightarrow{1/20}$	0.048/0.062
ξ_y	0.052	0.129/0.090	0.09/0.081
σ_z (mm)	4	6/7	6/5
I_{beam} (A)	2.6/1.1	1.64/1.19 $\xrightarrow{\times 2}$	3.6/2.6
$N_{bunches}$	5000	1584	2500
Luminosity ($10^{34} \text{cm}^{-2} \text{s}^{-1}$)	1.0	2.11 $\xrightarrow{\times 40}$	80


$$L = \frac{\gamma_{\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \left(\frac{I_{\pm} \xi_{y\pm}}{\beta_y^*} \right) \left(\frac{R_L}{R_{\xi_{y\pm}}} \right)$$

Cross sections at a B-factory


Physics process	Cross section [nb]	Selection Criteria	Reference
$\Upsilon(4S)$	1.110 ± 0.008	-	[2]
$u\bar{u}(\gamma)$	1.61	-	KKMC
$d\bar{d}(\gamma)$	0.40	-	KKMC
$s\bar{s}(\gamma)$	0.38	-	KKMC
$c\bar{c}(\gamma)$	1.30	-	KKMC
$e^+e^-(\gamma)$	300 ± 3 (MC stat.)	$10^\circ < \theta_e^* < 170^\circ$, $E_e^* > 0.15$ GeV	BABAYAGA.NLO
$e^+e^-(\gamma)$	74.4	$p_e > 0.5$ GeV/c and e in ECL	-
$\gamma\gamma(\gamma)$	4.99 ± 0.05 (MC stat.)	$10^\circ < \theta_\gamma^* < 170^\circ$, $E_\gamma^* > 0.15$ GeV	BABAYAGA.NLO
$\gamma\gamma(\gamma)$	3.30	$E_\gamma > 0.5$ GeV in ECL	-
$\mu^+\mu^-(\gamma)$	1.148	-	KKMC
$\mu^+\mu^-(\gamma)$	0.831	$p_\mu > 0.5$ GeV/c in CDC	-
$\mu^+\mu^-\gamma(\gamma)$	0.242	$p_\mu > 0.5$ GeV in CDC, $\geq 1 \gamma$ ($E_\gamma > 0.5$ GeV) in ECL	-
$\tau^+\tau^-(\gamma)$	0.919	-	KKMC
$\nu\bar{\nu}(\gamma)$	0.25×10^{-3}	-	KKMC
$e^+e^-e^+e^-$	39.7 ± 0.1 (MC stat.)	$W_{\ell\ell} > 0.5$ GeV/c ²	AAFH
$e^+e^-\mu^+\mu^-$	18.9 ± 0.1 (MC stat.)	$W_{\ell\ell} > 0.5$ GeV/c ²	AAFH

E. Kou, P. Urquijo et al.,
arXiv:1808.10567

Electromagnetic Calorimeter (ECL)




In barrel ECL, Belle II has **no projective cracks in ϕ** w.r.t. BaBar:
→ more hermetic
→ more efficient




Electromagnetic Calorimeter (ECL)


Energy resolution in Belle II barrel:



Material budget in front of ECL:




Beam background



Effects from beam background:

- degrades calorimeter resolution.
- radiation damage.
- pile-up and event size.
- physics background




BEAST: dedicated systems for continuous beam background measurement and monitoring!

See P. Lewis et al.: [10.1016/j.nima.2018.05.071](https://doi.org/10.1016/j.nima.2018.05.071)

Tracking performance


Tag & probe technique with events $e^+e^- \rightarrow \tau^+\tau^-$:

- 4 charged tracks expected;
- leptonID tags the event;
- 3 prongs provide the probe.



$$\epsilon = N_{\text{evts}}(4 \text{ tracks}) / [N_{\text{evts}}(4 \text{ tracks}) + N_{\text{evts}}(3 \text{ tracks})]$$

Track finding paper: arXiv:2003.12466
Performance plots: BELLE2-NOTE-PL-2020-014



Similar technique to evaluate **fake rate per track**:

$$r_{\text{fake}} = (0.97 \pm 0.34_{\text{stat}} \pm 0.06_{\text{syst}}) \%$$




Vertexing performance

D⁰ lifetime:

- measurement with 3 channels: $K^- \pi^+$, $K^- \pi^+$, $K^- \pi^+ \pi^- \pi^+$
- estimated vertex resolution: $\sim 40 \mu\text{m}$;

$$\rightarrow \tau(D^0) = (412.3 \pm 2.0_{\text{stat}}) \text{ fs}$$

(world average: $(410.1 \pm 1.5) \text{ fs}$)



B⁰ lifetime:

- smaller boost $\beta\gamma$: 0.42 (Belle) \rightarrow 0.28 (Belle II);
- average distance between B-mesons: $200 \mu\text{m} \rightarrow 130 \mu\text{m}$;
- hadronic channels: $B^0 \rightarrow D^{(*)-} \pi^+ / \rho^+$;
- estimated resolutions: $\Delta t \sim 1 \text{ ps} \rightarrow \Delta z \sim 80 \mu\text{m}$



$$\rightarrow \tau(B^0) = (1.48 \pm 0.28_{\text{stat}} \pm 0.06_{\text{syst}}) \text{ ps}$$

(world average: $(1.519 \pm 0.004) \text{ ps}$)

Particle identification: leptons

Several fully reconstructed channels are used:

- extract both efficiency & mis-ID probability **from data**;
- measured for various leptonID and angular acceptance.





$$\text{leptonID} = \mathcal{L}_{\text{lepton}} / (\mathcal{L}_e + \mathcal{L}_\mu + \mathcal{L}_\pi + \mathcal{L}_K + \mathcal{L}_p)$$

Particle identification: hadrons



Fully reconstructed channels are used:

- $D^{*+} \rightarrow D^0(\rightarrow K^-\pi^+)\pi^+$;
- slow π tags the D^0 flavour \rightarrow **K** and π **identified in data**.




$$K/\pi\text{-ID} = \mathcal{L}_{K/\pi} / (\mathcal{L}_\pi + \mathcal{L}_K)$$

Single Photon Trigger





L1 (hardware) trigger

Low Multiplicity Triggers



Measurement done with τ -pair events, 1x1 topology


Low Multiplicity Triggers



Measurement done with τ -pair events,
1x1 topology

L1 (hardware) trigger

A rule of thumb...



The masses of the mediator and of the DM candidates lead to **different type of searches**.



Dark Photon: invisible decay (signal)

Signal signature:

- a single, mono-chromatic, high-E photon (**ISR photon**)
- a bump in the recoil mass:

$$E_{\gamma} = \frac{s - m_{A'}^2}{2\sqrt{s}}$$


Needed a special **single photon trigger**
(not available in Belle, only ~10% of all data in BaBar)



Dark Photon: invisible decay (signal)

Discriminant variables:

E_{CMS} vs. polar angle of “single photon”




Signal signature:

peak in E_{CMS} (horizontal band)


Dark Photon: invisible decay (background)

Discriminant variables:

E_{CMS} vs. polar angle of "single photon"




$ee \rightarrow 2\gamma$ and 3γ
 1 γ in ECL 90° gap
 1 γ out of ECL acceptance




$ee \rightarrow eey$
 both electrons
 out of tracking acceptance

Signal signature:
 peak in E_{CMS} (horizontal band)





$ee \rightarrow 2\gamma$
 1 γ in ECL BWD or FWD gap



$ee \rightarrow 3\gamma$
 1 γ in ECL BWD gap
 1 γ out of ECL acceptance

Dark Photon: invisible decay




Single photon search: needed

a **special trigger logic**:

- ready for Belle II;
- not available in Belle;
- partially available in BaBar.

Dark Photon: leptonic decay




Look for a bump in the e^+e^- or $\mu^+\mu^-$ invariant mass over a (large) QED background

Belle II sensitivity is obtained by scaling the BaBar measurement:

- **expected better invariant mass resolution**
- **expected better triggers**

Recoil mass spectrum (after basic selections)




We found a deficit of data w.r.t. to MC (-35%)...

After having applied trigger and tracking corrections, in addition to a 65% correction, we obtain a very good agreement between data and MC.

NB: all the corrections are evaluated on independent control samples!

Suppression of the $\tau\tau$ background


The largest background component is due to $\tau\tau$ events: needed a special technique to suppress it!
→ Studied several variables, isolated the most discriminating ones between generated signal and background samples.



We optimized the selection in each recoil mass bin by choosing the best cuts that maximizes a given figure of merit
(hand-made multivariate approach).

$\tau\tau$ background suppressed

$\mu\mu$ final state



Suppression of the $\tau\tau$ background very effective up to 7 GeV (then, $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ events start to be dominant).


Signal efficiency between 3% and 5%.

No (local) anomalies observed... :(

And why not considering a LFV Z' ?

We considered only the $e-\mu$ coupling.
We considered only the invisible decay.

I. Galon et al. (2016), *arXiv:1610.08060*




Unfortunately, the model we were using showed some issues (too large width for the Z' , etc.).


**We decided to drop the signal model
and we opted for a model-independent search!**

LFV Z'

$e\mu$ final state



We can set limits on a coupling constant if a theoretical model is provided!



Axion-Like Particles (signal)

$$\tau_a \sim 1/g_{a\gamma\gamma}^2 m_a^3$$

For **resolved** case:

3 clusters with $E_{CM} > 0.25$ GeV



Peak in $\gamma\gamma$ mass spectrum

Three **resolved**,
high energetic
photons.



The searches for invisible and visible ALP decays veto this region.

Two of the
photons overlap
or **merge**.





ALP decays outside of the detector or decays into **invisible** particles: Single photon final state.

Axion-Like Particles: $a \rightarrow \gamma\gamma$

Search conducted with 445 pb⁻¹ of **2018 pilot run data**:


- 500 fits in sliding ranges with steps of half mass resolution;
- no excess observed (**largest local significance: 2.8 σ**).



ALPs: low-mass region

Belle II: ALPs below 200 MeV?



- ▶ For ALP masses below ~ 200 MeV, the decay photons are reconstructed as one ECL cluster even in offline analysis. Currently under study:
 - ▶ Untagged (electrons not seen) ALP fusion production has a much higher cross section and produces ALPs with less boost (difficult to trigger).
 - ▶ Shower shapes for merged cluster are different, MVA based reconstruction has better separation power (but events have to pass L1 trigger).
 - ▶ Pair conversion of one decay photon costs statistics, but yields a distinctive four particle final state.



Pro: resolved clusters

Con: very low energetic photons

Axion-like Particles from B decays



Izaguirre et al. (2017), *arXiv:1611.09355*