

The Belle II Experiment: status and prospects

Alberto Martini

**On behalf of the Belle II collaboration,
University & INFN Roma Tre**

**XIII International Conference on Beauty,
Charm and Hyperon Hadrons (BEACH)
Peniche, Portugal, 21 June 2018**



Belle II: second generation B-factory (I)

Main experiments at B-factories of the past:



- Belle (KEK Laboratory, Japan)
- BaBar (SLAC Laboratory, California)




Important results: confirmation of the CKM mechanism in the SM, CP violation observation in the B meson system etc..

Main problem:


statistics collected by KEKB and PEP-II colliders was not sufficient to analyse some rare decays, SM validations and other highly precise measurements



Belle II: usage of the improved collider **SuperKEKB**



Belle II: second generation B-factory (II)



By increasing the luminosity it is possible to investigate the SM through high precise measurements for highly suppressed, highly accurate or prohibited processes.



Improved detector performances




New Physics possibility!




Collider SuperKEKB

$$L = \frac{\gamma_{\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \frac{I_{\pm} \xi_{y\pm}}{\beta_{y\pm}^*} \frac{R_L}{R_{\xi_y}}$$



Collider SuperKEKB

$$L = \frac{\gamma_{\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \frac{I_{\pm} \xi_{y\pm}}{\beta_{y\pm}^*} \frac{R_L}{R_{\xi_y}}$$



KEKB

$$I_{e^+/e^-} = 1.64/1.19A$$


SuperKEKB

$$I_{e^+/e^-} = 3.6/2.6A$$



Collider SuperKEKB

$$L = \frac{\gamma_{\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \frac{I_{\pm} \xi_{y\pm}}{\beta_{y\pm}^*} \frac{R_L}{R_{\xi_y}}$$



KEKB

$$I_{e^+/e^-} = 1.64/1.19A$$

$$\beta_{y\ e^+/e^-}^* = 5.9/5.9mm$$

SuperKEKB


$$I_{e^+/e^-} = 3.6/2.6A$$

$$\beta_{y\ e^+/e^-}^* = 0.27/0.30mm$$



Collider SuperKEKB

$$L = \frac{\gamma_{\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \frac{I_{\pm} \xi_{y\pm}}{\beta_{y\pm}^*} \frac{R_L}{R_{\xi_y}}$$



KEKB



SuperKEKB

$$I_{e^+/e^-} = 1.64/1.19 A$$

$$I_{e^+/e^-} = 3.6/2.6 A$$

$$\beta_{y\ e^+/e^-}^* = 5.9/5.9 mm$$

$$\beta_{y\ e^+/e^-}^* = 0.27/0.30 mm$$




Beam section at the interaction point:
 ~42 nm in y
 ~6 μm in x

Nano-beam scheme



Collider SuperKEKB

$$L = \frac{\gamma_{\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \frac{I_{\pm} \xi_{y\pm}}{\beta_{y\pm}^*} \frac{R_L}{R_{\xi_y}}$$



KEKB



SuperKEKB

$$I_{e^+/e^-} = 1.64/1.19 A$$

$$I_{e^+/e^-} = 3.6/2.6 A$$

$$\beta_{y\ e^+/e^-}^* = 5.9/5.9 mm$$

$$\beta_{y\ e^+/e^-}^* = 0.27/0.30 mm$$




Beam section at the interaction point:
 ~42 nm in y
 ~6 μm in x

Nano-beam scheme




Expected improvement of **integrated luminosity** of a factor ~50 w.r.t. Belle: **50 ab⁻¹**

Belle II detector (I)




Belle II detector (II)

Increased luminosity → Higher occupancy, pile-up issues, more background hits
Higher trigger and DAQ rate
Radiation damage increase




Improvements with respect to Belle:

- Signal readout speed and waveform fit in the e.m. calorimeter;
- K_S reconstruction efficiency (+30%);
- K/π separation (wrong ID probability reduced by a factor ~ 2.5);
- Primary and secondary vertices reconstruction (resolution x2).




Belle II detector (II)

Increased luminosity → Higher occupancy, pile-up issues, more background hits
Higher trigger and DAQ rate
Radiation damage increase




Improvements with respect to Belle:

- Signal readout speed and waveform fit in the e.m. calorimeter;
- K_S reconstruction efficiency (+30%);
- K/π separation (wrong ID probability reduced by a factor ~ 2.5);
- Primary and secondary **vertices reconstruction** (resolution x2).



Belle II status and plan



Phase 1: SuperKEKB commissioning & background estimation

Completed

Phase 2: Collision runs with the detector installed partially, without the vertex detector → first physics data!

Ongoing and almost finished!

Phase 3: Data taken with the whole detector installed

February 2019




Belle II first event displays

First hadronic event!



Phase 2 first hadronic physics event!




First beautiful hadronic event!




Belle II first results

First kaon mass reconstruction plot



Vertical beam size σ_y^* as a function of the date





Instant luminosity reached so far by SuperKEKB $\sim 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$




Physics program

- Precision measurement of the Unitarity Triangle angles and CKM matrix elements.





- New CP violation sources through time-dependent/integrated asymmetry measurements.
- Flavour Changing Neutral Current (FCNC) studies
- Search for Lepton Flavour Violation (LFV)
- Dark sector investigation
- Hadronic spectroscopy and quarkonium studies

details...




Belle II exclusive advantages

- Coherent production of two B mesons from $\Upsilon(4S)$ resonance
- “Clean” environment w.r.t. experiments using hadronic machine:
 - Large data samples with B, D and τ with low background
 - Analysis of decays with missing energy
- Good reconstruction efficiency and resolution for neutral particles as γ , K , π^0



Belle II exclusive advantages

- Coherent production of two B mesons from $\Upsilon(4S)$ resonance
- “Clean” environment w.r.t. experiments using hadronic machine:
 - Large data samples with B, D and τ with low background
 - Analysis of decays with missing energy
- Good reconstruction efficiency and resolution for neutral particles as γ , K , π^0




Full Event Interpretation (FEI)


Full reconstruction of one B meson decay (B_{tag}) \rightarrow determination of the flavour of the other B (B_{sig}) and isolation of particles coming from the signal side \rightarrow (large advantages in analysis with missing energy/mass)

Reconstruction efficiency:

- 1.5% semi-leptonic tag: more efficient but less precise
- 0.3% hadronic tag: less efficient but more precise



Lepton flavour violation in τ decays



SM prediction:

$$BF(\tau \rightarrow l\gamma) \sim 10^{-40}$$


Studied only at
the B-factory

LHCb not competitive

details..

Signal side

$$\tau^- \rightarrow \mu^- \gamma$$




Tag side

$$\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$$

Belle II expectations:

Improvement of ≈ 2
order of magnitude
w.r.t. the actual limits




Flavour anomalies in $R(D^*)$ and $R(D)$

Observables:

$$R(D^*) = \frac{BF(B \rightarrow D^* \tau \nu)}{BF(B \rightarrow D^* \mu \nu)} \stackrel{SM}{=} 0.252 \pm 0.003$$

4.1 σ SM disagreement





Flavour anomalies in $R(D^*)$ and $R(D)$

Observables:

$$R(D^*) = \frac{BF(B \rightarrow D^* \tau \nu)}{BF(B \rightarrow D^* \mu \nu)} \stackrel{SM}{=} 0.252 \pm 0.003$$

4.1 σ SM disagreement



It could be explained through the existence of **charged Higgs** or other New Physics models


More details in Mario Merola's talk



Dark sector

Minimal model introducing the dark interaction comprises:


- A' : dark photon. Boson mediator of the dark interaction with mass $m_{A'}$ and spin 1
- ϵ : coupling parameter. It indicates the coupling intensity between the dark photon and the SM photon




Dark sector

Minimal model introducing the dark interaction comprises:


- A' : dark photon. Boson mediator of the dark interaction with mass $m_{A'}$ and spin 1
- ϵ : coupling parameter. It indicates the coupling intensity between the dark photon and the SM photon
- $\chi_{1,2}$: dark matter particles



A' decay into invisible states



At the moment, this analysis can be performed by Belle II thanks to the **single photon** dedicated trigger.




Possible phase 2 analysis $\mathcal{L}^{\text{int}} = 20 \text{ fb}^{-1}$




Dark sector

Minimal model introducing the dark interaction comprises:


- A' : dark photon. Boson mediator of the dark interaction with mass $m_{A'}$ and spin 1
- ϵ : coupling parameter. It indicates the coupling intensity between the dark photon and the SM photon
- $\chi_{1,2}$: dark matter particles



A' decay into invisible states



At the moment, this analysis can be performed by Belle II thanks to the **single photon** dedicated trigger.



Possible phase 2 analysis $\mathcal{L}^{\text{int}} = 20 \text{ fb}^{-1}$



Conclusions


- New collider SuperKEKB $\rightarrow \mathcal{L}^{\text{int}} = 50 \text{ ab}^{-1}$ before 2026
- Improved detector performances: good neutral particle reconstruction, resonances, decay vertices and events with high missing energy.
- Fundamental physics studies: CKM matrix, CPV, LFV, FCNC, dark sector.
- Installation and insertion of the detector: 11 April 2017
- Current status, phase 2: first data analysed without vertex detector.
- Phase 3: data taking will start in February 2019 with the whole detector installed.




Belle II insertion
11 April 2017



First physics event!!
25 April 2018



Emergency slides!!



Accessible channels and σ at Belle II

Number of particles produced, assuming 100% of beam on each resonance.

Channel	Belle	BaBar	Belle II (per year)
$B\bar{B} \Upsilon(4S)$	7.7×10^8	4.8×10^8	1.1×10^{10}
$B_s^{(*)}\bar{B}_s^{(*)}$	7.0×10^6	—	6.0×10^8
$\Upsilon(1S)$	1.0×10^8		1.8×10^{11}
$\Upsilon(2S)$	1.7×10^8	0.9×10^7	7.0×10^{10}
$\Upsilon(3S)$	1.0×10^7	1.0×10^8	3.7×10^{10}
$\Upsilon(5S)$	3.6×10^7	—	3.0×10^9
$\tau\tau$	1.0×10^9	0.6×10^9	1.0×10^{10}


Process	Cross section (nb)	Rate (Hz)
$\Upsilon(4S) \rightarrow B\bar{B}$	1.2	960
$e^+e^- \rightarrow$ continuum	2.8	2200
$\mu^+\mu^-$	0.8	640
$\tau^+\tau^-$	0.8	640
Bhabha ($\theta_{\text{lab}} \geq 17^\circ$)	44	350 ^a
$\gamma\gamma$ ($\theta_{\text{lab}} \geq 17^\circ$)	2.4	19 ^a
2γ processes ^b	~ 80	~ 15000
Total	~ 130	~ 20000

Total event rate: 20 KHz allowed, thanks to improved detector system performances.

^a Rate is pre-scaled by a factor 1/100


^b $\theta_{\text{lab}} \geq 17^\circ, p_t \geq 0.1\text{GeV}/c$ Alberto Martini - BEACH2018 21/06/2018

CM boost



$$E_{e^+} \simeq 4 \text{ GeV}$$

$$E_{e^-} \simeq 7 \text{ GeV}$$



- E_{CM} at the resonance of $\Upsilon(4S)$

- Center of mass boost:
 $\beta\gamma = 0.28$

Symmetric beams:

$$\beta\gamma \simeq 0.06 \longrightarrow \Delta r \simeq 30 \mu\text{m}$$



**Decay vertex can not
be resolve.**

Asymmetric beams:

$$\beta\gamma \simeq 0.28 \longrightarrow \Delta z = \beta\gamma \cdot c \cdot \tau \simeq 130 \mu\text{m}$$



**Decay vertex can be
resolved.**

Belle CM boost: $\beta\gamma = 0.425 \rightarrow$ more separation but less luminosity.

Lepton flavour violation in τ decays (I)

Ratios of BFs of τ LFV decays allow to discriminate NP models!


	SUSY+GUT (SUSY+Seesaw)	Higgs mediated	Little Higgs	Non-universal Z'
$\frac{\mathcal{B}(\tau \rightarrow \mu\mu\mu)}{\mathcal{B}(\tau \rightarrow \mu\gamma)}$	$\sim 2 \cdot 10^{-3}$	0.06-0.1	0.4-2.3	~ 16
$\frac{\mathcal{B}(\tau \rightarrow \mu ee)}{\mathcal{B}(\tau \rightarrow \mu\gamma)}$	$\sim 1 \cdot 10^{-2}$	$\sim 1 \cdot 10^{-2}$	0.3-1.6	~ 16
$\mathcal{B}(\tau \rightarrow \mu\gamma)_{max}$	$< 10^{-7}$	$< 10^{-10}$	$< 10^{-10}$	$< 10^{-9}$



$$B \longrightarrow K^{*0} \nu \bar{\nu}$$

SM prediction box diagram + penguin:

$$BF_{SM}(B \longrightarrow K^{*0} \nu \bar{\nu}) = (9.48 \pm 1.10) \cdot 10^{-6}$$



Upper limit measured at Belle:

$$BF_{Belle}(B \longrightarrow K^{*0} \nu \bar{\nu}) < 5.5 \cdot 10^{-5}$$

Could lead to New Physics!

SM validity can be proved at 5σ through the decay rate measurement using the whole Belle II statistic: $\mathcal{L}^{\text{int}} = 50 \text{ ab}^{-1}$





$$B^0 \longrightarrow K^{*0} \mu^+ \mu^-$$

FCNC: b → s transitions

Possible New Physics

Angular analysis (using P_5') chosen to reduce theoretical uncertainties



Previous analysis done by the Belle experiment shows a discrepancy of P_5' parameter within a certain q^2 range of $\sim 2.6\sigma$ in the SM → comparable results with LHCb analysis.





$$B^0 \longrightarrow K^{*0} \mu^+ \mu^-$$

FCNC: $b \rightarrow s$ transitions

Possible New Physics




$$R_{K^{(*)}} = BR(B \rightarrow K^{(*)} \mu\mu) / (B \rightarrow K^{(*)} ee)^{SM} \simeq 1$$

Belle II projection sensitivity on R_{K^*} :



Belle II experiment contribution will be crucial for that measurement!

Hadronic spectroscopy



S. Godfrey and K. Moats, Bottomonium mesons and strategies for their observation, Phys. Rev. D 92, 054034 (2015)
 S. Godfrey and N. Isgur, Mesons in a relativized quark model with chromodynamics, Phys. Rev. D 32, 189 (1985).

Many intermediate states have been observed and many others have not

QCD knowledge at low energies is needed to interpretate possible New Physics signals.

back..

Complementary to LHCb

Observable	Expected th. accuracy	Expected exp. uncertainty	Facility
CKM matrix			
$ V_{us} [K \rightarrow \pi \ell \nu]$	**	0.1%	<i>K</i> -factory
$ V_{cb} [B \rightarrow X_c \ell \nu]$	**	1%	Belle II
$ V_{ub} [B_d \rightarrow \pi \ell \nu]$	*	4%	Belle II
$\sin(2\phi_1) [c\bar{c}K_S^0]$	***	$8 \cdot 10^{-3}$	Belle II/LHCb
ϕ_2		1.5°	Belle II
ϕ_3	***	3°	LHCb
CPV			
$S(B_s \rightarrow \psi\phi)$	**	0.01	LHCb
$S(B_s \rightarrow \phi\phi)$	**	0.05	LHCb
$S(B_d \rightarrow \phi K)$	***	0.05	Belle II/LHCb
$S(B_d \rightarrow \eta' K)$	***	0.02	Belle II
$S(B_d \rightarrow K^*(\rightarrow K_S^0 \pi^0) \gamma))$	***	0.03	Belle II
$S(B_s \rightarrow \phi \gamma))$	***	0.05	LHCb
$S(B_d \rightarrow \rho \gamma))$		0.15	Belle II
A_{SL}^d	***	0.001	LHCb
A_{SL}^s	***	0.001	LHCb
$A_{CP}(B_d \rightarrow s \gamma)$	*	0.005	Belle II
rare decays			
$B(B \rightarrow \tau \nu)$	**	3%	Belle II
$B(B \rightarrow D \tau \nu)$		3%	Belle II
$B(B_d \rightarrow \mu \nu)$	**	6%	Belle II
$B(B_s \rightarrow \mu \mu)$	***	10%	LHCb
zero of $A_{FB}(B \rightarrow K^* \mu \mu)$	**	0.05	LHCb
$B(B \rightarrow K^{(*)} \nu \nu)$	***	30%	Belle II
$B(B \rightarrow s \gamma)$		4%	Belle II
$B(B_s \rightarrow \gamma \gamma)$		$0.25 \cdot 10^{-6}$	Belle II (with 5 ab^{-1})
$B(K \rightarrow \pi \nu \nu)$	**	10%	<i>K</i> -factory
$B(K \rightarrow e \pi \nu) / B(K \rightarrow \mu \pi \nu)$	***	0.1%	<i>K</i> -factory
charm and τ			
$B(\tau \rightarrow \mu \gamma)$	***	$3 \cdot 10^{-9}$	Belle II
$ q/p _D$	***	0.03	Belle II
$arg(q/p)_D$	***	1.5°	Belle II

Both LHCb and Belle II are needed to cover all the precision flavour physics aspects.

LHCb:

- Decay channels with charged particles in the final state.


Belle II:

- Decay channels with some neutrinos or neutral particles in the final state;
- Inclusive decays;
- Decay channels involving long lived particles: K_S & K_L .

back..

B. Golob, KEK FF Workshop, Feb. 2012

SuperKEKB luminosity projection



[back...](#)

