

BELLE2-NOTE-PL-2019-008
DRAFT Version 2.0
September 29, 2019

Approved plots of untagged $B \rightarrow D^*l\nu_l$ decays with 5.15 fb^{-1} Proc9 and bucket7 early Phase III Data

The Belle II Collaboration

Abstract

We report the plots approved for Beauty 2019 conference for untagged $B \rightarrow D^*l\nu_l$ result using 5.15 fb^{-1} of early phase III DATA. Details are in the internal note BELLE2-NOTE-PH-2019-019.




FIG. 1: The Maximum likelihood fit to $\cos \theta_{BY} = \frac{2E_B^* E_Y^* - M_B^2 - m_Y^2}{2p_B^* p_Y^*}$ and $m_{\text{miss}}^2 = \left(\frac{P_{ee}}{2} - P_Y^*\right)^2$ distributions of untagged $\bar{B}^0 \rightarrow D^{*+} l^- \bar{\nu}_l$ (charge conjugate modes taken in to account) candidates using 5.15 fb^{-1} of collision Data, where E_Y^* , p_Y^* , P_Y^* , and m_Y are the center-of-mass (CM) energy, three momentum, four momentum and invariant mass of the $D^* l$ system, P_{ee} is the four momentum of the beam particles, M_B is the nominal B mass, and E_B^* , p_B^* are the CM energy and momentum of the B , inferred from the CM machine energy. For correctly reconstructed B candidates, ignoring detector resolution effects and the spread in machine energy, θ_{BY} is the CM angle between the B and $D^* l$ momenta. Here Data are shown with points with error bars with different components overlaid for $\bar{B}^0 \rightarrow D^{*+} e^- \bar{\nu}_e$ (top) and $\bar{B}^0 \rightarrow D^{*+} \mu^- \bar{\nu}_\mu$ (bottom) channels. D^0 candidates are reconstructed from $K^- \pi^+$ pairs, selected without particle identification requirements, within the invariant mass range $1.85 \text{ GeV}/c^2 < m_{K\pi} < 1.88 \text{ GeV}/c^2$. D^{*+} candidates are reconstructed from a D^0 candidate and a π^+ candidate track, with the invariant-mass difference between D^{*+} and D^0 candidates in the range $0.144 \text{ GeV}/c^2 < \Delta m < 0.148 \text{ GeV}/c^2$. The CM momentum of the D^{*+} candidate is required to satisfy $p_{D^{*+}}^* < 2.5 \text{ GeV}/c$. Continuum $e^+ e^- \rightarrow q\bar{q}$ background is suppressed with the Fox-Wolfram moment ratio $R2 < 0.25$. The CM momentum of the lepton candidate is required to be in the range $1.2 \text{ GeV}/c < p_l^* < 2.4 \text{ GeV}/c$. Electron and muon candidates are selected with requirements on the combined variables, $\text{electronID} > 0.85$ and $\text{muonID} > 0.9$ respectively. We observe O(1100) $\bar{B}^0 \rightarrow D^{*+} e^- \bar{\nu}_e$ and O(1200) $\bar{B}^0 \rightarrow D^{*+} \mu^- \bar{\nu}_\mu$ events.




FIG. 2: Projection of ΔM and p_l^* in the signal enhanced region of $|m_{\text{miss}}^2| < 1.5 \text{ GeV}/c$. Here Data are shown with points with error bars with different components overlaid for $\bar{B}^0 \rightarrow D^{*+} e^- \bar{\nu}_e$ (top) and $\bar{B}^0 \rightarrow D^{*+} \mu^- \bar{\nu}_\mu$ (bottom) channels.