



BELLE2-NOTE-PL-2019-018

Version 2.1

August 5, 2019

## Approved plots for $J/\psi \rightarrow \ell^+\ell^-$ in Proc9

D. Ferlewicz,<sup>\*</sup> M. Milesi,<sup>†</sup> and P. Urquijo<sup>‡</sup>

*School of Physics, University of Melbourne, Victoria, Australia*

### Abstract

Approved plots for the dilepton yields of  $J/\psi \rightarrow \ell^+\ell^-$  for the analysis documented in BELLE2-NOTE-PH-2019-050.

---

<sup>\*</sup>Electronic address: [daniel.ferlewicz@unimelb.edu.au](mailto:daniel.ferlewicz@unimelb.edu.au)

<sup>†</sup>Electronic address: [marco.milesi@unimelb.edu.au](mailto:marco.milesi@unimelb.edu.au)

<sup>‡</sup>Electronic address: [purquijo@unimelb.edu.au](mailto:purquijo@unimelb.edu.au)




FIG. 1: The dielectron invariant mass of  $J/\psi \rightarrow e^-e^-$  candidates for an integrated luminosity of  $2.62 \text{ fb}^{-1}$  using the `basf2` software release `release-03-02-02` on the `hlt_hadron` skim. This data set includes Phase 3 physics runs only, excluding runs 916 – 1005 and 1216 – 1371 from experiment 7. The selection criteria are as follows:  $|dr| < 2.0 \text{ cm}$ ,  $|dz| < 5.0 \text{ cm}$ ,  $p_{\text{lab}} > 0.1 \text{ GeV}/c$  and `electronID`  $> 0.95$  for each electron candidate. A vertex fit using `TreeFitter` was applied, selecting candidates with a p-value  $> 0.001$ . A bremsstrahlung correction was applied by adding the momentum and cluster energy of a photon with  $E < 1.0 \text{ GeV}$  within a  $5^\circ$  cone of the electron candidate.

A Crystal Ball function summed with a Bifurcated Gaussian is used to model the signal and a first order polynomial is used to model the background.




FIG. 2: The dimuon invariant mass of  $J/\psi \rightarrow \mu^+\mu^-$  candidates for an integrated luminosity of  $2.62 \text{ fb}^{-1}$  using the same environment and track selection as Fig. 1, but with  $\text{muonID} > 0.95$  for each muon candidate.

A Gaussian function summed with a Bifurcated Gaussian is used to model the signal and a first order polynomial is used to model the background.




FIG. 3: The dielectron invariant mass of  $J/\psi \rightarrow e^-e^-$  candidates for an integrated luminosity of  $2.62 \text{ fb}^{-1}$  using the same environment and track selection as the Fig. 1, but with further selection criteria applied to  $J/\psi$  candidates in  $B\bar{B}$  events. The momentum of the reconstructed  $J/\psi$  candidate in the  $\Upsilon(4S)$  frame is required to be below  $2.0 \text{ GeV}/c$  and the ratio between the (event-based) second-order and zeroth-order Fox-Wolfram moment,  $R_2$ , is below  $0.4$ .




FIG. 4: The dimuon invariant mass of  $J/\psi \rightarrow \mu^+\mu^-$  candidates for an integrated luminosity of  $2.62 \text{ fb}^{-1}$  using the same environment and track selection as Fig. 2, with extra selection criteria applied to isolate  $J/\psi$  candidates in  $B\bar{B}$  events, listed in Fig. 3.