

Radiative and electroweak penguin decay at Belle II

Keisuke Yoshihara, on behalf of the Belle II Collaboration

Department of Physics and Astronomy, Iowa State University, Ames, IA 50011, USA

E-mail: Keisuke.Yoshihara@cern.ch

Flavor changing neutral current b to $s(d)$ decay, particularly radiative and electroweak penguin processes, will continue to be of great interest as a probe to search for beyond the Standard Model physics. Lately there have been several anomalies observed in the b to s transition at the LHCb experiment in an angular observable, P_5' and lepton universality measurements. The Belle II experiment will take an important role providing independent cross checks for these anomalies. This article describes how the Belle II physics program can be complementary to that of LHCb, and the Belle II ongoing studies and prospects on radiative and electroweak penguin processes in early and final dataset of 50 ab^{-1} .

BEAUTY2020
21-24 September 2020
Kashiwa, Japan (online)

1. Introduction

The Standard Model of particle physics (SM) is a well established theoretical framework. However, there still remain unanswered questions such as baryon asymmetry in universe or the origin of dark matter, naively indicating the necessity of physics beyond the SM (BSM). As B -meson decays via flavor changing neutral current (FCNC) are forbidden at tree-level in the SM, these decays only proceed through higher-order loop diagrams. Radiative and electroweak penguin processes shown in Figure 1 are typical example of such a B -meson FCNC decay.




Figure 1: The leading order Feynman diagrams for FCNC decays: (a) radiative penguin and (b) electroweak penguin process in the SM.

These penguin decays may receive contribution from BSM at a comparable size. Moreover, as the BSM appears as off-shell particles, these decays can provide a unique opportunity to explore higher mass scale compared to searches involving direct production. Several recent measurements in $b \rightarrow sl^+\ell^-$ decays are in tension with the SM predictions. For example, a measurement of the decay-angular distributions in the decay $B^0 \rightarrow K^{*0}\mu^+\mu^-$ performed by LHCb, shows more than 2.5σ discrepancy with the SM for the observable P'_5 [1, 2]. Measurements of the ratio of branching fractions of the decays $B \rightarrow K^{(*)}\mu^+\mu^-$ and $B^+ \rightarrow K^{(*)}e^+e^-$, referred to as R_K and R_K^* , are also found to be about 2.5σ inconsistent with the lepton flavor universality [3, 4]. The Belle II experiment will be able to provide independent cross-check for these measurements, possibly finding a hint for BSM physics. Prospects for various measurements in the context of radiative and electroweak penguin decays are described in the following sections.

2. SuperKEKB and Belle II detector

SuperKEKB [5] is an e^+e^- collider, located at KEK in Tsukuba, Japan, with asymmetric beam energy of 4 GeV for positrons and 7 GeV for electrons such that the collision energy is equal to the energy of the $\Upsilon(4S)$ resonance and $B\bar{B}$ pairs get boosted in forward direction. SuperKEKB was designed and constructed as an upgrade of KEKB accelerator complex, aiming to collect total integrated luminosity of 50 ab^{-1} with the maximum peak luminosity of $6 \times 10^{35}\text{ cm}^{-2}\text{ s}^{-1}$, which is approximately 30 times higher than the recorded peak luminosity of KEKB. This dataset will contain approximately $50 \times 10^9 B\bar{B}$ events. The beam interaction point is surrounded by Belle II detector [6],

a multipurpose detector with a 4π solid angle composed of various sub-detector systems; pixel and silicon strip layers (PXD and SVD), central drift chamber (CDC), particle identification counters (TOP and ARICH), electromagnetic calorimeter (ECL), and K_L -muon detector (KLM). The Belle II detector was also designed and constructed as an upgrade of Belle detector to be capable of reconstructing vertices of B -mesons precisely (vertex resolution of $\ll 50\mu\text{m}$ is required) while coping with the higher luminosity as well as the higher beam-induced background. Belle II detector started taking physics data since 2019 and the total integrated luminosity of 74 fb^{-1} was collected to date (September 2020).

3. Analysis strategy

3.1 Theoretical framework

Effective Hamiltonian is a general expression to describe a full theory at low energy (μ) and it has been used widely to interpret results in a model independent way. The Hamiltonian can be factorized into local operators O_i with Wilson coefficients C_i as follows:

$$\mathcal{H}_{eff} \sim \sum_i C_i(\mu) O_i(\mu) \quad (1)$$

There are in total 24 operators in the full expression. However, only three operators are relevant in $b \rightarrow s$ process. Radiative penguin process is involved as O_7 and electroweak penguin as O_9 (vector) and O_{10} (axial-vector). The BSM contribution is expected to modify SM contributions or introduce new operators in the form of $\Delta\mathcal{H} \sim (C_{BSM}/\Lambda_{BSM})O_{BSM}$ where Λ_{BSM} is the cut-off energy of new physics. Global fit with various different measurements is able to obtain model-independent constraints on C_i and hence constrain BSM models.

3.2 B -meson reconstruction

As B -mesons are pair-produced, Belle II will be able to perform inclusive measurements where the hadronic part of the decay is not specified. The inclusive measurements are complementary to the exclusive ones as they tend to be less affected by uncertainties due to hadronic form-factors. For the reconstruction of $B \rightarrow X_s \gamma$ decays, for example, there are two “inclusive” methods used and they are often referred to as (1) sum-of-exclusive method and (2) fully-inclusive method. In the sum-of-exclusive method, the hadronic system is reconstructed from exclusive decays containing a kaon and pion(s). Hadronic candidates are then combined with a hard photon to reconstruct B -meson candidates. The B -meson is referred to as B_{sig} . In the fully-inclusive method, the other B -meson referred to as B_{tag} is also reconstructed to improve the S/B ratio. B_{tag} is classified further into three types of tag; (1) a fully reconstructed hadronic final state (hadronic tagging), (2) a fully reconstructed semi-leptonic decay (semi-leptonic tagging), and (3) only an energetic lepton (leptonic tagging) from the B -meson decay. These reconstruction methods have their own pros and cons providing access to different observables. Full event interpretation (FEI) [7] is a new reconstruction algorithm developed for Belle II employing machine learning technique and the reconstruction efficiency of B_{tag} is expected to improve approximately by a factor of two in comparison to the efficiency obtained by the Belle algorithm.




Figure 2: A schematic of a hadronically tagged decay of the B -meson (B_{tag}) where $B \rightarrow D^0(\rightarrow K\pi)\pi$ is reconstructed in this example. Several hadronic decay chains with high branching fractions or efficiencies are included as B_{tag} . For the signal (B_{sig}), the B -meson decay being targeted is reconstructed. The signal can include neutrinos as missing energy or an inclusive decay (X_S) where all possible hadrons are considered.

4. Radiative penguin decay

$B \rightarrow X_S \gamma$ offers important constraints through C_7 on many BSM scenarios such as supersymmetry and models with extended Higgs sectors. The prompt photons are selected as isolated clusters in ECL that are not matched to any charged tracks. In order to reduce contaminations from asymmetric $\pi^0/\eta \rightarrow \gamma\gamma$ decays, each photon candidate is paired with all other photons in the event. The B -mesons are then reconstructed as described in Section 3.2.




Figure 3: The Belle II sensitivity projection ?? for $BF(B \rightarrow X_S \gamma)$ (left) and $(\Delta)\mathcal{A}_{CP}(B \rightarrow X\gamma)$ (right) as a function of integrated luminosity. The green line in BF is discontinued assuming the improvement of hadronic tagging in the Belle II experiment.

For the inclusive measurement of branching fraction (BF), the sensitivity of previous measurements has been limited by different sources of uncertainties. For the leptonic tagging, the uncertainty is dominated by the systematic uncertainty arising from neutral hadrons faking photons. The uncertainty is expected to be reduced significantly at Belle II by exploiting the cluster shape information in ECL. While for the hadronic tagging, the uncertainty was statistically dominated due to the low reconstruction efficiency of B -mesons. In case of the sum-of-exclusive method, the dominant uncertainty arises from hadronization of X_S such as fragmentation and missing decay

modes. The sensitivity projection for $BF(B \rightarrow X_s\gamma)$ is shown in Figure 3. $BF(B \rightarrow X_s\gamma)$ will be measured in 6% precision with the early dataset of 5 ab^{-1} and in 3% precision with full dataset of 50 ab^{-1} . In addition to the BF measurements, other important observables such as CP asymmetry (\mathcal{A}_{CP}), isospin asymmetry (Δ_{0+}), and difference of CP asymmetry ($\Delta\mathcal{A}_{CP}$) as defined in Equation 2 and 3 can be measured at Belle II.

$$\mathcal{A}_{CP} = \frac{\Gamma[\bar{B} \rightarrow X_s\gamma] - \Gamma[B \rightarrow X_s\bar{\gamma}]}{\Gamma[\bar{B} \rightarrow X_s\gamma] + \Gamma[B \rightarrow X_s\bar{\gamma}]}, \quad \Delta_{0+} = \frac{\Gamma[B^0 \rightarrow X_s\gamma] - \Gamma[B^\pm \rightarrow X_s\gamma]}{\Gamma[B^0 \rightarrow X_s\gamma] + \Gamma[B^\pm \rightarrow X_s\gamma]}, \quad (2)$$

$$\Delta\mathcal{A}_{CP} = \mathcal{A}_{CP}[B^\pm \rightarrow X_s^\pm\gamma] - \mathcal{A}_{CP}[B^0 \rightarrow X_s^0\gamma], \quad (3)$$

In these measurements, experimental systematic uncertainties can be reduced along with theoretical uncertainty on hadronic form-factors. Figure 3 also shows sensitivity projection for these measurements.

5. Electroweak penguin decay

$B \rightarrow X_s\ell^+\ell^-$, where ℓ is either e or μ , also provides important information on the quark flavor sector that is complementary to the $B \rightarrow X_s\gamma$ decay, and constrains C_9 and C_{10} . Electrons are selected as an isolated track that is matched to a cluster in ECL with a large p/E . Charged tracks that leave cluster hits in KLM are selected as muons. Belle II is capable of performing inclusive and exclusive measurements of the $B \rightarrow X_s\ell^+\ell^-$ decay.




Figure 4: The Belle II sensitivity projection ?? for $BF(B \rightarrow X_s\ell^+\ell^-)$ (left) and \mathcal{A}_{FB} (right) in three different regions of squared invariant mass of the lepton pair.

In the inclusive mode, hadronic uncertainties can be significantly suppressed compared to the exclusive mode, and hence higher experimental sensitivity is expected. The branching fraction of $B \rightarrow X_s\ell^+\ell^-$ decay can be measured in three regions of squared invariant mass of the lepton pair. Figure 4 shows the sensitivity projection for the branching fraction measurement as a function of integrated luminosity. In addition to the BF measurements, the forward-backward asymmetry

(\mathcal{A}_{FB}) and a difference of \mathcal{A}_{FB} ($\Delta_{CP}(\mathcal{A}_{FB}) \equiv \mathcal{A}_{FB}^{\bar{B}} - \mathcal{A}_{FB}^B$) can be also measured as a function of the dilepton invariant mass at Belle II. The sensitivity projection for the \mathcal{A}_{FB} measurement is also shown in Figure 4. The \mathcal{A}_{FB} measurement is expected to obtain the most stringent constraint on C_9 and C_{10} as theoretical and experimental systematic uncertainties are largely canceled. The uncertainty on the BF measurements will be dominated by theoretical uncertainties at 15 ab^{-1} while the \mathcal{A}_{FB} measurement will be statistically dominated even at 50 ab^{-1} . Furthermore, \mathcal{A}_{CP} can be measured at Belle II. The \mathcal{A}_{CP} of the $B \rightarrow X_s \ell^+ \ell^-$ decay can be a sensitive probe for new sources of CP violation. Test of lepton flavor universality is also an important subject. The $R(K)$ and $R(K^*)$ anomalies can be cross checked in low- and high- q^2 regime with the higher sensitivity than previous measurements by taking an advantage of the high reconstruction efficiency and purity for both electrons and muons. In addition, Belle II will perform an inclusive R_{X_s} measurement which is expected to help largely cancel systematic uncertainties. Moreover, angular observables such as P'_5 can be measured in low- and high- q^2 regime and another lepton universality test can be performed with Q_5 ($\equiv P'_5 e - P'_5 \mu$) observable. The expected sensitivity in R_{X_s} and P'_5 (and Q_5) as a function of the integrated luminosity is shown in Figure 5.




Figure 5: The Belle II sensitivity projection ?? for R_{X_s} (left) and P'_5 (and Q_5) (right) as a function of integrated luminosity.

6. $B \rightarrow K^{(*)}\nu\bar{\nu}$ decay

$B \rightarrow K^{(*)}\nu\bar{\nu}$ is another probe that is sensitive to C_9 and C_{10} . It is theoretically and experimentally cleaner than $B \rightarrow X_s \ell^+ \ell^-$ because of no photon mediated contribution. The $B \rightarrow K^{(*)}\nu\bar{\nu}$ decay has never been observed in previous experiments. Belle II is expected to be the first experiment to observe the decay mode with 5 ab^{-1} assuming the SM rate. With the full dataset, the branching fraction can be measured in about 10% accuracy. Since two neutrinos are present in the final state, the other B -meson needs to be fully reconstructed. The FEI algorithm described above is expected to take an essential role in this measurement. The fraction of longitudinal polarization (F_L) is also known to be sensitive to BSM signature and the detailed study can follow with the full dataset.

7. Conclusion

Rare radiative and electroweak processes have drawn a lot of attention from the high energy physics community because of several anomalies observed in previous measurements (mainly performed by LHCb) in $b \rightarrow s$ transition. The SuperKEKB / Belle II experiment started physics data-taking in 2019 and 74 fb^{-1} has been collected to date (September 2020). Belle II will have access to decay modes and measurements that are likely challenging at LHCb, such as the $B \rightarrow K^{(*)}\nu\bar{\nu}$ decay and various inclusive measurements, and will be able to provide an independent verification to these anomalies, possibly finding a hint for BSM in the near future.

References

- [1] LHCb Collaboration, “Angular analysis of the $B^0 \rightarrow K^{*0}\mu^+\mu^-$ decay using 3 fb^{-1} of integrated luminosity”, JHEP02 (2016)104.
- [2] LHCb Collaboration, “Measurement of CP-Averaged Observables in the $B \rightarrow K^0\mu^+\mu^-$ Decay”, PRL125.011802(2020).
- [3] LHCb Collaboration, “Test of lepton universality with $B^0 \rightarrow K^{*0}\ell^+\ell^-$ decays”, JHEP08(2017)055.
- [4] LHCb Collaboration, “Search for lepton-universality violation in $B^+ \rightarrow K^+\ell^+\ell^-$ decay”, PRL122.191801(2019).
- [5] Y. Onishi et al, “Accelerator design at SuperKEKB”, <https://doi.org/10.1093/ptep/pts083>
- [6] Belle II Collaboration, “Belle II Technical Design Report”, arXiv:1011.0352.
- [7] T. Keck et al., “The Full Event Interpretation - An exclusive tagging algorithm for the Belle II experiment”, Comput. Softw. Big Sci. 3 (2019) , arXiv:1807.08680 [hep-ex].
- [8] Belle II Collaboration, “Belle II Physics Book”, <https://arxiv.org/abs/1808.10567>.