



BELLE2-NOTE-PL-2021-001

Approved Plots: Υ dipion transitions in Early Phase 3




FIG. 1: Plot of $M(\mu\mu)$ versus $M(\pi^+\pi^-\mu^+\mu^-) - M(\mu^+\mu^-)$ for $\Upsilon \rightarrow \mu^+\mu^-$ dipion decays. The clusters of points represent signal transitions.

Belle II Preliminary

$\int L dt = 72 \text{ fb}^{-1}$




FIG. 2: Plot of $M(\pi^+\pi^-\mu^+\mu^-) - M(\mu^+\mu^-) + m(\Upsilon(1S))$ with a requirement of $|M(\mu\mu) - m(\Upsilon(1S))| < 75 \text{ MeV}/c^2$, where $m(\Upsilon(1S))$ represents the nominal $\Upsilon(1S)$ mass.

Belle II Preliminary

$\int L dt = 72 \text{ fb}^{-1}$




FIG. 3: Plot of $M(\pi^+\pi^-\mu^+\mu^-) - M(\mu^+\mu^-) + m(\Upsilon(2S))$ with a requirement of $|M(\mu\mu) - m(\Upsilon(2S))| < 75 \text{ MeV}/c^2$, where $m(\Upsilon(2S))$ represents the nominal $\Upsilon(2S)$ mass.