



QUARKONIUM/ QCD RESULTS AT BELLE II

M.C. Chang

Fu Jen Catholic University, TW

On behalf of the [Belle II](#) Collaboration

3 July , 2023



Beauty 2023

BELLE II EXPERIMENT




Asymmetric e^+e^- collider in Tsukuba, Japan

Nano-beam interaction
point



Tunable E_{cm} around $\Upsilon(4S)$ mass
 $L = 6.0 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ (ultimate)
 $L = 4.7 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ (Belle II)





SuperKEKB

Belle II detector

QUARKONIA



= $c\bar{c}$ and $b\bar{b}$ mesons (or charmonia and bottomonia)



EXOTIC HADRONS

XYZ states:


lots of them in charmonium

bottomonium analogues: Y_b , Z_b , Z'_b

what are they?




which partons compose them?




tetraquark




pentaquark



hybrid



hadronic molecule



glueball

THE BELLE LEGACY



Belle@KEKB (B-factory) → optimized for

$$e^+e^- \rightarrow \Upsilon(4S) \rightarrow B\bar{B}$$

However

$\Upsilon(5S)$: discovery of $hb(1,2P)$, $\eta_b(2S)$, $Z_b(10610,10650)$


[PR D91 072003, PRL 109 232002]

exotic states and anomalous $\pi\pi$ transition widths

Process	Partial width
$\Upsilon(10860) \rightarrow \Upsilon(3S)\pi^+\pi^-$	$(0.59 \pm 0.04 \pm 0.09) \text{ MeV}$
$\Upsilon(10860) \rightarrow \Upsilon(2S)\pi^+\pi^-$	$(0.85 \pm 0.07 \pm 0.09) \text{ MeV}$
$\Upsilon(10860) \rightarrow \Upsilon(1S)\pi^+\pi^-$	$(0.52_{-0.17}^{+0.20} \pm 0.10) \text{ MeV}$
$\Upsilon(3S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	$(8.9 \pm 0.8) \times 10^{-4} \text{ MeV}$

Energy scan data: $\Upsilon(10753)$ aka Y_b

rise in hadronic transition cross sections (resonance)




DISCOVERY OF $\Upsilon(10753)$



Observed in the $e^+ e^- \rightarrow \Upsilon(nS) \pi^+ \pi^-$ ($n = 1, 2, 3$) cross section energy dependence by Belle (JHEP 10 (2019) 220):


	$\Upsilon(10860)$	$\Upsilon(11020)$	New structure
M (MeV/c ²)	$10885.3 \pm 1.5^{+2.2}_{-0.9}$	$11000.0^{+4.0}_{-4.5} {}^{+1.0}_{-1.3}$	$10752.7 \pm 5.9^{+0.7}_{-1.1}$
Γ (MeV)	$36.6^{+4.5}_{-3.9} {}^{+0.5}_{-1.1}$	$23.8^{+8.0}_{-6.8} {}^{+0.7}_{-1.8}$	$35.5^{+17.6}_{-11.3} {}^{+3.9}_{-3.3}$



A dip in $\sigma(e^+e^- \rightarrow b\bar{b})$ can be described by $\Upsilon(10753)$
CPC 44, 8, 083001
(2020)

WHAT'S THE NATURE OF $\Upsilon(10753)$

- Far from the thresholds;





- Mass does not match $\Upsilon(3D)$ theoretical predictions, and D-wave states are not seen in e^+e^- collisions;
- $\Upsilon(4S) - \Upsilon(3D)$ mixing can be enhanced due to hadron loops.

- Tetraquark state:
[CPC 43, 12, 123102 \(2019\)](#),
[PLB, 802, 135217 \(2020\)](#),
- Hadronic molecule with a small admixture of a bottomonium:
[PRD 103, 074507 \(2021\)](#)
- Hybrid state:
[PRD 99, 1, 014017 \(2019\)](#)
- Conventional $b\bar{b}$ state:
[EPJC 80, 1, 59 \(2020\)](#)
[PLB 803, 135340 \(2020\)](#)
[PRD 102, 1, 014036 \(2020\)](#)
[PRD 101, 1, 014020 \(2020\)](#)
[PRD 104, 034036 \(2021\)](#)
[PRD 105, 074007 \(2022\)](#)
[PRD 106, 094013 \(2022\)](#)
[EPJC 137, 357 \(2022\)](#)

ABOVE $\Upsilon(4S)$: NOV. 2021 ENERGY SCAN



$\Upsilon(10753)$ state was observed in the $e^+ e^- \rightarrow \Upsilon(nS) \pi^+ \pi^-$ ($n = 1, 2, 3$) cross section energy dependence by Belle (JHEP 10 (2019) 220).




- ❑ 19 fb^{-1} scan around $\Upsilon(10753)$ was collected in November 2021;
- ❑ Belle II collected the data in the gaps between Belle energy scan points;
- ❑ The point with highest statistic (9.8 fb^{-1}) is near $\Upsilon(10753)$ peak;


STUDY OF $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$

- ❑ $\sigma(e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)})$ energy dependence show complicated spectra, that hard to describe with resonance shapes;
- ❑ Rescattering and opening of the various $B\bar{B}$ thresholds cause oscillatory behaviour due to the **coupled-channel effect**;
- ❑ **Coupled-channel approach** is necessary to study $\sigma(e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)})$ shape;

JHEP 06 (2021) 137



GLOBAL PHENOMENOLOGICAL ANALYSIS



Data:

- ☐ Two-body exclusive cross sections $\sigma(e^+e^- \rightarrow B_{(s)}^{(*)} \bar{B}_{(s)}^{(*)})$;
- ☐ Three-body exclusive cross sections $\sigma(e^+e^- \rightarrow \Upsilon(nS) \pi^+ \pi^-)$, $n = 1, 2, 3$;
 $\sigma(e^+e^- \rightarrow h_b(mP) \pi^+ \pi^-)$, $m = 1, 2$;
- ☐ Combined Belle and BaBar R_b measurement;
- ☐ **Use coupled-channel approach.**
- ☐ **Poles:** $\Upsilon(4S)$, $\Upsilon(10753)$, $\Upsilon(5S)$ and $\Upsilon(6S)$
- ☐ **Results:** pole positions (mass and width), branching fractions, dependence of scattering amplitudes on energy.

PRD 106 (2022) 9, 094013

TWO BELLE II RESULTS WILL BE PRESENTED



More data is necessary

- ❑ To study $Y(10753)$ nature;
- ❑ Improve accuracy below $Y(5S)$;




Perform energy scan at the Belle II experiment.



Two Belle II results will be presented:

- ❑ $e^+e^- \rightarrow \omega \chi_{bJ}(1P)$ and $X_b \rightarrow \omega Y(1S)$
- ❑ $e^+e^- \rightarrow B\bar{B}, B\bar{B}^*$ and $B^*\bar{B}^*$



Search for $e^+e^- \rightarrow \omega \chi_{bJ}(1P)$ and $X_b \rightarrow \omega \Upsilon(1S)$

PRL 130 091902 (2023)




Motivation to search for $Y(10753) \rightarrow \omega \chi_{bJ}(1P)$

Theory:


- Mixed $Y(4S) - Y(3D)$ state: $\omega \chi_{bJ}$ could be enhanced (PRD **104**, 034036 (2021)).

Charmonium sector:

- Similar to $Y(10753)$ structure $Y(4220)$ was observed in $e^+e^- \rightarrow J/\psi \pi^+\pi^-$ cross section dependence by BES III (PRL **118**, 092001 (2017)).
- $Y(4220)$ peak was observed in $\gamma X(3872)$ and $\omega \chi_{c0}$ final states by BES III (PRL, **122**, 232002 (2019), PRD **99**, 091103(R) (2019)).
- We can expect $Y(10753)$ to decay into $\gamma[X_b \rightarrow \omega Y(1S)]$ and $\omega \chi_{bJ}$ final states.



Observation of $\Upsilon(10753) \rightarrow \omega \chi_{bJ}(1P)$




PRL **130**, 091902 (2023)

Channel	\sqrt{s} (GeV)	N^{sig}	$\Sigma(\sigma)$	σ_B (pb)
$e^+e^- \rightarrow \omega \chi_{b0}$	10.701	< 3.0	-	< 16.6
$e^+e^- \rightarrow \omega \chi_{b1}$		< 3.9	-	< 1.2
$e^+e^- \rightarrow \omega \chi_{b2}$		< 4.0	-	< 2.5
$e^+e^- \rightarrow \omega \chi_{b0}$	10.745	< 12.0	0.5	< 11.3
$e^+e^- \rightarrow \omega \chi_{b1}$		$68.9^{+13.7}_{-13.5}$	5.9	$3.6^{+0.7}_{-0.7} \pm 0.5$
$e^+e^- \rightarrow \omega \chi_{b2}$		$27.6^{+11.6}_{-10.0}$	3.1	$2.8^{+1.2}_{-1.0} \pm 0.4$
$e^+e^- \rightarrow \omega \chi_{b0}$	10.805	< 9.9	1.2	< 11.4
$e^+e^- \rightarrow \omega \chi_{b1}$		$15.0^{+6.8}_{-6.2}$	2.7	< 1.7
$e^+e^- \rightarrow \omega \chi_{b2}$		$3.3^{+5.3}_{-3.8}$	0.8	< 1.6


11σ

4.5σ

2D fit to $M(\gamma \Upsilon(1S))$ and $M(\pi^+ \pi^- \pi^0)$:



Observation of $\Upsilon(10753) \rightarrow \omega \chi_{bJ}(1P)$




PRL 130, 091902 (2023)

Fit with coherent sum of PHSP and BW.

$$\left| \sqrt{\Phi_2(\sqrt{s})} + \frac{\sqrt{12\pi\Gamma_{ee}\mathcal{B}_f\Gamma}}{s - M^2 - iM\Gamma} \sqrt{\frac{\Phi_2(\sqrt{s})}{\Phi_2(M)}} e^{i\phi} \right|^2$$

M and Γ are fixed to 10752.7 MeV and 35.5 MeV

- ☐ Confirms $\Upsilon(10753)$ state;
- ☐ No peak at $\Upsilon(5S)$;
- ☐ $\sigma(\chi_{b1}\omega)/\sigma(\chi_{b2}\omega) \sim 1$;
- ☐ $\sigma(\chi_{b1}\omega)/\Upsilon(2S)\pi^+\pi^- \sim 1.5$;



$\Gamma_{ee}\mathcal{B}_f$	Solution I	Solution II
	(constructive interference)	(destructive interference)
$\Gamma_{ee}\mathcal{B}_f(\Upsilon(10753) \rightarrow \omega\chi_{b1})$	$(0.63 \pm 0.39 \pm 0.20)$ eV	$(2.01 \pm 0.38 \pm 0.46)$ eV
$\Gamma_{ee}\mathcal{B}_f(\Upsilon(10753) \rightarrow \omega\chi_{b2})$	$(0.53 \pm 0.46 \pm 0.15)$ eV	$(1.32 \pm 0.44 \pm 0.55)$ eV

Discussion

Previously Belle measured $\sigma(e^+e^- \rightarrow \chi_{bJ}(1P)\omega)$ at $\sqrt{s} = 10.867$ GeV (PRL **113** (2014) 14, 142001);

- Y(5S) and Y(10753) have same quantum numbers and similar masses, but there is a difference:

$$\frac{\sigma(e^+e^- \rightarrow \chi_{bJ}(1P)\omega)}{\sigma(e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-)} \sim \begin{cases} \sim 1.5 \text{ at } \sqrt{s} = 10.745 \text{ GeV} \\ \sim 0.15 \text{ at } \sqrt{s} = 10.867 \text{ GeV} \end{cases}$$


- Order of magnitude difference is observed for this ratio at Y(5S) and Y(10753)



It indicates the difference in the internal structures of these two states.



Search for $\Upsilon(10753) \rightarrow \gamma X_b [\rightarrow \omega \Upsilon(1S)]$



- ❑ No evidence of X_b (partner of $X(3872)$ in bottomonium) signal;
- ❑ Only $\omega \chi_{bJ}(1P)$ reflections are seen;


$\Upsilon(1S) (\rightarrow e^+e^-/\mu^+\mu^-)$ PRL 130, 091902 (2023)

- ❑ Upper limits on cross sections are set for $M(X_b) \in [10.45; 10.65]$ GeV;

$$\sigma_{X_b}^{\text{UL}} = \sigma_B^{\text{UL}}(e^+e^- \rightarrow \gamma X_b) \mathcal{B}(X_b \rightarrow \omega \Upsilon(1S))$$

\sqrt{s} (GeV)	M_{X_b} (GeV)	$\sigma_{X_b}^{\text{UL}}$ (pb)
10.653	10.59	< 0.55
10.701	10.45	< 0.84
10.745	10.45	< 0.14
10.805	10.53	< 0.47

Fit to $M[\omega \Upsilon(1S)]$



Energy dependence of the $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$ cross section

Energy dependence of the $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$ cross section

Previous Belle analysis: [JHEP 06 \(2021\), 137](#)


- ❑ One B meson is fully reconstructed using hadronic channels;
- ❑ $B^* \rightarrow B\gamma$ decays are not reconstructed;

$$\Delta E = E_B - E_{\text{cm}}/2$$





$$\Delta E' = \Delta E + M_{\text{bc}} - m_B$$

- ❑ $|\Delta E'| < 18 \text{ MeV}$;
- ❑ Signal is identified using M_{bc} :

$$M_{\text{bc}} = \sqrt{E_{\text{cm}}^2/4 - p_B^2}$$



M_{bc} FIT AT SCAN ENERGIES





- ❑ Good description of the M_{bc} in data;
- ❑ Contribution of $\Upsilon(4S) \rightarrow B\bar{B}$ production via ISR is visible well described by the fit;
- ❑ $E=10.653$ GeV sharp cut of the data at right edge \Rightarrow fast rise of $B^*\bar{B}^*$ near threshold;

ENERGY DEPENDENCE OF THE CROSS SECTIONS

Simultaneous fit to:



- Exclusive cross sections measured by in this work and previous Belle study ([JHEP 06 \(2021\), 137](#));

- Total cross section ([CPC 44, 8, 083001 \(2020\)](#))



COMPARISON OF $\sigma_{b\bar{b}}$ and $\sigma_{B\bar{B}} + \sigma_{B\bar{B}^*} + \sigma_{B^*\bar{B}^*}$

- Good agreement at low energies;
- Difference at higher energy is due to $B_s^{(*)}\bar{B}_s^{(*)}$, multi-body $B^{(*)}\bar{B}^{(*)}\pi(\pi)$ and bottomonia;




DISCUSSION

New measurement complements previous Belle result:

- ❑ Solid curve – combined Belle + Belle II data fit
- ❑ Dashed curve – Belle data fit only

$\sigma(e^+e^- \rightarrow B^*\bar{B}^*)$ rises rapidly above $B^*\bar{B}^*$ threshold:

- ❑ Similar behaviour was seen for $D^*\bar{D}^*$ cross section ([PRD 97, 012002 \(2018\)](#));
- ❑ **Possible interpretation:** resonance or bound state ($b\bar{b}$ or $B^*\bar{B}^*$) near threshold ([MPL A 21, 2779 \(2006\)](#));
- ❑ Also explains a narrow dip in $\sigma(e^+e^- \rightarrow B\bar{B}^*)$ near $B^*\bar{B}^*$ threshold by destructive interference between $e^+e^- \rightarrow B\bar{B}^*$ and $e^+e^- \rightarrow B^*\bar{B}^* \rightarrow B\bar{B}^*$;
- ❑ $\Upsilon \pi^+ \pi^-$ and $h_b \eta$ final states could also be enhanced ([PRD 87, 094033 \(2013\)](#)).



CONCLUSION

Observation of $e^+e^- \rightarrow \omega\chi_{bJ}(1P)$ at $\sqrt{s} = 10.75$ GeV

- $\sigma[e^+e^- \rightarrow \omega\chi_{bJ}(1P)]$ has a peak at 10.75 GeV
- Confirmation of $Y(10753)$ and observation of its new decay channel;

Energy dependence of $e^+e^- \rightarrow B\bar{B}, B\bar{B}^*$ and $B^*\bar{B}^*$

- Confirmation of “oscillatory” behavior, improvement of the accuracy;
- Rapid rise of $\sigma(e^+e^- \rightarrow B^*\bar{B}^*)$ above threshold - signal of molecular $B^*\bar{B}^*$ state?

Scan above $Y(4S)$ gives an opportunity for a lot of unique studies:

- $Y(10753)$ decays to different final states. Study of its properties;
- Energy dependence of the various final states production;

Golden Modes

$$e^+e^- \rightarrow \pi^+\pi^-\Upsilon(pS)(\rightarrow \ell^+\ell^-)$$

$B\bar{B}$ decomposition Preliminary result

$$\pi^+\pi^- \text{ Dalitz}$$

$$Y_b \rightarrow \omega\eta_b(1S)$$

$Y_b \rightarrow \omega\chi_{bJ}(1P)$ [PRL 130, 091902 \(2023\)](#)

Silver Modes

$$Y_b \rightarrow \pi^+\pi^- X \text{ (inclusive)}$$

$$Y_b \rightarrow \eta X \text{ (inclusive)}$$

$$Y_b \rightarrow \eta\Upsilon(1S, 2S)(\rightarrow \ell^+\ell^-)$$

$$Y_b \rightarrow \eta'\Upsilon(1S)(\rightarrow \ell^+\ell^-)$$

$$Y_b \rightarrow \Upsilon(1S) \text{ (inclusive)}$$

Bronze Modes

$$Y_b \rightarrow \gamma X_b$$

$$Y_b \rightarrow \pi^0\pi^0\Upsilon(pS)(\rightarrow \ell^+\ell^-)$$

$$Y_b \rightarrow KK(\phi)\Upsilon(pS)(\rightarrow \ell^+\ell^-)$$

$$Y_b \rightarrow \pi^0\pi^0 X \text{ (inclusive)}$$

$$Y_b \rightarrow \pi^0 X \text{ (incl. or excl.)}$$

...