

## The Study of Radiative $D_s$ Decays

---

**N Sushree Ipsita,<sup>a,\*</sup> Vishal Bhardwaj<sup>b</sup> and Anjan Giri<sup>a</sup>**

*(On behalf of Belle collaboration)*

<sup>a</sup>*Department of Physics*

*Indian Institute of Technology, Hyderabad, India*

<sup>b</sup>*Department of Physics*

*IISER Mohali, India*

*E-mail: [ph19resch02005@iith.ac.in](mailto:ph19resch02005@iith.ac.in)*

The study of weak radiative decays of charmed mesons is still in its developing stage. In the Standard Model (SM), the physics of charm mesons is not generally expected to have New Physics (NP) discovery potential. The weak decays of  $D$  mesons are also difficult to investigate due to the strong final-state interactions related to QCD.  $c \rightarrow u\gamma$  decays can be affected by some contributions coming from the non-minimal supersymmetry, which is a NP scenario.  $R_{\rho/\omega}$  could be violated already in the SM framework, while a similar relation for  $D_s^+$  radiative decays offers a much better test for  $c \rightarrow u\gamma$ . Here, we present a sensitivity study of the radiative charm decays  $D_s^+ \rightarrow \rho^+\gamma$  and  $D_s^+ \rightarrow K^{*+}\gamma$  with data collected by the Belle experiment.

*16th International Conference on Heavy Quarks and Leptons - HQL2023  
November 28 to December 2, 2023  
Mumbai, India*

---

\*Speaker

## 1. Introduction

In the Standard Model (SM), the physics of charmed mesons is not generally expected to have New Physics (NP) discovery potential because of the relevant CKM matrix [1] elements  $V_{cs}$  and  $V_{cd}$  are well known and the CP asymmetries and  $D^0 - \bar{D}^0$  oscillations are small. Further, the weak decays of  $D$  mesons are difficult to investigate due to the strong final-state interactions related to QCD. However, it has been pointed out that the oscillations and  $c \rightarrow u\gamma$  decays might have some contributions coming from the non-minimal supersymmetry (the NP scenario). Therefore, one can search for NP using  $c \rightarrow u\gamma$  transitions. It was suggested that the NP would result in a deviation from  $R_{\rho/\omega}$  [2]:

$$R_{\rho/\omega} \equiv \frac{\Gamma(D^0 \rightarrow \rho^0/\omega\gamma)}{\Gamma(D^0 \rightarrow \bar{K}^{*0}\gamma)} = \frac{\tan^2\theta_c}{2} \quad (1)$$

In order to find the best mode to test  $c \rightarrow u\gamma$  decay, the ratios between various Cabibbo-suppressed and Cabibbo-allowed radiative decays of charmed mesons are calculated as predicted by the SM. It has been noticed that Eq. (1) could be violated already in the SM framework because of a large, unknown correction within the SM, while a similar relation for  $D_s^+$  radiative decays offers a much better test to search for a signal of NP, as this ratio is less sensitive to the SM parameters [3].

$$R_K \equiv \frac{\Gamma(D_s^+ \rightarrow K^{*+}\gamma)}{\Gamma(D_s^+ \rightarrow \rho^+\gamma)} = \tan^2\theta_c \quad (2)$$

Radiative  $D_s$  decays, such as  $D_s^+ \rightarrow K^{*+}\gamma$  and  $D_s^+ \rightarrow \rho^+\gamma$ , have not been observed yet. The theoretical analysis of the  $D \rightarrow V\gamma$  transitions was done using a model that combines heavy quark effective theory and the chiral Lagrangian approach and includes symmetry breaking [4]. In addition to the s-channel annihilation and t-channel W exchange, there is a long-distance penguin-like  $c \rightarrow u\gamma$  contribution in the Cabibbo-suppressed modes. Its magnitude is determined by the size of symmetry breaking, which was calculated with a vector dominance approach. Although smaller in magnitude, the penguin-like contribution would lead to sizable effects in case of cancellations among the other contributions to the amplitude. Thus, it may invalidate suggested tests beyond the SM effects in these decays. Figure 1 shows the Feynman diagram of  $c \rightarrow u\gamma$  transition [5]. This




Figure 1: Feynman diagram

model predicts a range of values for the branching ratios predicted for the various  $D \rightarrow V\gamma$  modes, as shown in table 1. Here, we present the first experimental study of these modes.


Decay Mode	Branching Fraction
$D_s^+ \rightarrow \rho^+ \gamma$ [4]	$(3-5) \times 10^{-4}$
$D_s^+ \rightarrow K^{*+} \gamma$ [4]	$(2.1-3.2) \times 10^{-5}$

**Table 1:** Summary of the expected value of the branching fraction.

## 2. Sample Selection


The selection of signal candidates is optimized using simulated samples that have been generated with the EvtGen [6] and Geant packages [7]. We have reconstructed  $D_s^+$  from  $D_s^+ \rightarrow \rho^+ \gamma$  and  $D_s^+ \rightarrow K^{*+} \gamma$ , where  $\rho^+ \rightarrow \pi^+ \pi^0$  and  $K^{*+} \rightarrow K_s^0 \pi^+$ , respectively. The kinematic variable that distinguishes the signal from the background is  $\Delta M$ , where  $\Delta M$  is the difference between the reconstructed mass of  $D_s^{*+}$  and  $D_s^+$  ( $\Delta M \equiv M(D_s^{*+}) - M(D_s^+)$ ). To reduce the combinatorial background, we keep only those candidates that satisfy the criteria:  $0.08 \text{ GeV}/c^2 < \Delta M < 0.20 \text{ GeV}/c^2$ . A  $\pi^0$  veto has been implemented to get rid of the huge background coming from  $\pi^0$  decays. We have performed background MC study in which the continuum background is found to be dominant. We employ multivariate analysis (MVA) using the FastBDT package to get rid of uds background, processes such as  $e^+ e^- \rightarrow u\bar{u}, d\bar{d}, s\bar{s}$ . After applying a cut greater than 0.4 (0.5) on MVA for  $D_s^+ \rightarrow \rho^+ \gamma$  ( $D_s^+ \rightarrow K^{*+} \gamma$ ) decay mode, there is a rejection of 65% (76%) of uds background at the cost of 10% (24%) of signal loss. The reconstruction efficiency is found to be 0.5% (3.1%) for the  $D_s^+ \rightarrow \rho^+ \gamma$  ( $D_s^+ \rightarrow K^{*+} \gamma$ ), respectively. For the  $D_s^+ \rightarrow \rho^+ \gamma$  decay mode, the peaking background is mostly coming from the  $D_s^+ \rightarrow \rho^+ \eta$ , and for the  $D_s^+ \rightarrow K^{*+} \gamma$ , it is mostly coming from the  $D^0 \rightarrow K^0 \pi^0$  and  $D^0 \rightarrow K^0 \eta$  decay modes.

## 3. Control Sample Study

**Figure 2:** Fitted distribution of  $\Delta M$  for (a)  $D_s^+ \rightarrow \rho^+ \eta$ , (b)  $D^0 \rightarrow K_s^0 \pi^0$  and (c)  $D^0 \rightarrow K_s^0 \eta$  decay modes, respectively.

We utilize the peaking backgrounds  $D_s^+ \rightarrow \rho^+ [\eta \rightarrow \gamma\gamma]$ ,  $D^{*0} \rightarrow [D^0 \rightarrow K_s^0 \eta] \gamma$  and  $D^{*0} \rightarrow [D^0 \rightarrow K_s^0 \pi^0] \gamma$  as our control sample to verify the signal extraction procedure and to calibrate possible discrepancies in the signal resolution between data and simulation. Figure 2 shows the 1D unbinned maximum likelihood fit on  $\Delta M$  for (a)  $D_s^+ \rightarrow \rho^+ \eta$  and (b)  $D^0 \rightarrow K_s^0 \pi^0$  and (c)  $D^0 \rightarrow K_s^0 \eta$  decay modes, respectively.

#### 50 4. Signal Extraction



**Figure 3:** Fitted distribution of  $\Delta M$  for (a)  $D_s^+ \rightarrow \rho^+\gamma$  and (b)  $D_s^+ \rightarrow K^{*+}\gamma$  decay modes, respectively.

51 We have performed 1D unbinned maximum likelihood fit on  $\Delta M$  for both (a)  $D_s^+ \rightarrow \rho^+\gamma$  and  
 52 (b)  $D_s^+ \rightarrow K^{*+}\gamma$  decay modes, as shown in Figure 3. For  $D_s^+ \rightarrow \rho^+\gamma$  case, the signal is modeled  
 53 with the sum of two bifurcated Gaussian with a common mean. The peaking background is modeled  
 54 with the sum of two bifurcated Gaussian, combinatorial background with third-order Chebyshev  
 55 Polynomial. For  $D_s^+ \rightarrow K^{*+}\gamma$  mode, the signal is modeled with the sum of two bifurcated Gaussian.  
 56 The peaking backgrounds are modeled with the sum of a Gaussian and a bifurcated Gaussian. The  
 57 combinatorial background is modeled with third-order Chebyshev Polynomial. All the signal and  
 58 peaking background parameters are fixed except the Chebyshev Polynomial.

#### 59 5. Preliminary Results and Outlook

60 We are expecting 300-400 (20-30) events for  $D_s^+ \rightarrow \rho^+\gamma$  ( $D_s^+ \rightarrow K^{*+}\gamma$ ) decay modes using  
 61  $10^{-4}$  ( $10^{-5}$ ) branching fraction corresponding to an integrated luminosity of  $921 \text{ fb}^{-1}$ .

#### 62 References

- 63 [1] M. Kobayashi and T. Maskawa, *CP-Violation in the Renormalizable Theory of Weak*  
 64 *Interaction*, *Progress of Theoretical Physics* **49** (1973) 652.
- 65 [2] I.I. Bigi, *Weak decays of charm hadrons: The next lesson on qcd – and possibly more!*, 1993.
- 66 [3] B. Bajc *et al.*,  *$c \rightarrow u\gamma$  in cabibbo suppressed d meson radiative weak decays*, *Phys. Rev. D* **54**  
 67 (1996) 5883.
- 68 [4] S. Fajfer and P. Singer, *Long distance  $c \rightarrow u\gamma$  effects in weak radiative decays of d mesons*,  
 69 *Phys. Rev. D* **56** (1997) 4302.
- 70 [5] S. Fajfer *et al.*, *Long distance contributions in  $d \rightarrow v\gamma$  decays*, *Eur. Phys. J. C* **6** (1999) 471.
- 71 [6] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001)  
 72 152.
- 73 [7] R. Brun *et al.*, *GEANT3*, *CERN-DD-EE-84-1* (1987) .