

Charm physics prospects at the Belle II experiment

Longke LI (李龙科)

On behalf of the Belle II Collaboration

Institute of High Energy Physics, CAS, China (IHEP)



21st Particles and Nuclei International Conference
PANIC 2017, Sep. 1-5, Beijing

Outline

- 1 Belle II at SuperKEKB
- 2 D^0 - \bar{D}^0 mixing and CP violation
 - Formalism and experiment status
 - Time-dependent measurements
 - Time-integrated CP asymmetry
- 3 Rare or leptonic charm decays
 - Rare radiative decays with one photon
 - Rare radiative decays with two photons
 - (Semi-) Leptonic charm decays
- 4 ROE method of D^0 flavor tag
- 5 Summary



Outline


- 1 Belle II at SuperKEKB
- 2 D^0 - \bar{D}^0 mixing and CP violation
 - Formalism and experiment status
 - Time-dependent measurements
 - Time-integrated CP asymmetry
- 3 Rare or leptonic charm decays
 - Rare radiative decays with one photon
 - Rare radiative decays with two photons
 - (Semi-) Leptonic charm decays
- 4 ROE method of D^0 flavor tag
- 5 Summary



SuperKEKB

- ▶ KEKB \Rightarrow SuperKEKB (with "Nano-Beam" scheme)


more info, see [Jake's report](#) on 3rd Sep



$$\text{luminosity: } \mathcal{L} = \frac{\gamma_{\pm}}{2e r_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \frac{I_{\pm} \zeta_{y\pm}}{\beta_{y\pm}^*} \left(\frac{R_L}{R_{\zeta_y}} \right)$$

	E (GeV)	I (A)	β_y^* (mm)	Lumin.
	LER/HER	LER/HER	LER/HER	($cm^{-2}s^{-1}$)
KEKB	3.5/8.0	1.64/1.19	5.9/5.9	2.1×10^{34}
SuperKEKB	4.0/7.0	3.60/2.60	0.27/0.31	80×10^{34}
	$\beta\gamma \sim 2/3$	$\times 2$	$\times 20$	$\times 40$

- ▶ Target integrated luminosity for Belle II at SuperKEKB: $\int_t \mathcal{L} dt = 50 \text{ ab}^{-1}$




First physics run will start in 2018
Each 1 ab^{-1} experimental data provides

- $\sim 1.1 \times 10^9 B\bar{B} \Rightarrow$ a super **B-factory**;
- $\sim 1.3 \times 10^9 c\bar{c} \Rightarrow$ a super **charm factory**;
- $\sim 0.9 \times 10^9 \tau^+\tau^- \Rightarrow$ a super **τ factory**;
- wide effective $E_{c.m.}=[0.5-10]$ GeV via ISR process.



Belle II Detector



Impacts on Charm WG

- ☺ 50× larger dataset collected by Belle II
- ☺ Better D^0 proper time resolution (improved IP and secondary vertex resolution)
- ☺ Better reconstruction efficiency with improved tracking efficiency, eg: $D^{*+} \rightarrow D^0 \pi_s^+$ etc.
- ☺ Better particle identification capabilities, eg: $D^0 \rightarrow K^+ \pi^-$ etc.
- ☺ ...



Outline

- 1 Belle II at SuperKEKB
- 2 D^0 - \bar{D}^0 mixing and CP violation
 - Formalism and experiment status
 - Time-dependent measurements
 - Time-integrated CP asymmetry
- 3 Rare or leptonic charm decays
 - Rare radiative decays with one photon
 - Rare radiative decays with two photons
 - (Semi-) Leptonic charm decays
- 4 ROE method of D^0 flavor tag
- 5 Summary



Formalism of D^0 - \bar{D}^0 mixing and CP violation

- Open-flavor neutral meson transforms to anti-meson:

$$K^0 \Leftrightarrow \bar{K}^0, B_d^0 \Leftrightarrow \bar{B}_d^0, B_s^0 \Leftrightarrow \bar{B}_s^0, D^0 \Leftrightarrow \bar{D}^0$$


- Flavor eigenstate ($|D^0\rangle, |\bar{D}^0\rangle$) \neq mass eigenstate $|D_{1,2}\rangle$ with $M_{1,2}$ and $\Gamma_{1,2}$

$$|D_{1,2}\rangle \equiv p|D^0\rangle \pm q|\bar{D}^0\rangle \quad (\text{CPT: } p^2+q^2=1)$$

- Mixing parameters: $\mathbf{x} \equiv 2 \frac{M_1 - M_2}{\Gamma_1 + \Gamma_2}, \quad \mathbf{y} \equiv \frac{\Gamma_1 - \Gamma_2}{\Gamma_1 + \Gamma_2}$

- Unique system: only up-type meson for mixing

- Standard Model(SM) predicts: $\sim \mathcal{O}(1\%)$

(1) short distance ($< 0.1\%$)(2) long distance ($\sim 1\%$)

- Precise measurement of x, y : effectively limit the New Physics(NP) modes; and search for NP, eg: $|x| \gg |y|$

- Three types of **C**harged-conjugated-**P**arity combined symmetry **V**iolation (**CPV**):

$$A_{CP}^f = \frac{\Gamma(D \rightarrow f) - \Gamma(\bar{D} \rightarrow \bar{f})}{\Gamma(D \rightarrow f) + \Gamma(\bar{D} \rightarrow \bar{f})} = a_d^f + a_m^f + a_i^f$$

- a_d^f : (direct CPV) CPV in decay $|\bar{A}_f/A_f| \neq 1$

$$\left| \begin{array}{c} p^0 \\ \swarrow \searrow \\ \bullet \\ \swarrow \searrow \\ f \end{array} \right|^2 \neq \left| \begin{array}{c} \bar{p}^0 \\ \swarrow \searrow \\ \bullet \\ \swarrow \searrow \\ \bar{f} \end{array} \right|^2$$

- a_m^f : CPV in mixing with $r_m = |q/p| \neq 1$

$$\left| \begin{array}{c} p^0 \quad \bar{p}^0 \\ \swarrow \quad \swarrow \\ \bullet \\ \swarrow \quad \swarrow \\ \bar{f} \end{array} \right|^2 \neq \left| \begin{array}{c} \bar{p}^0 \quad p^0 \\ \swarrow \quad \swarrow \\ \bullet \\ \swarrow \quad \swarrow \\ f \end{array} \right|^2$$


- a_i^f : CPV in interference with $\arg(q/p) \neq 0$

$$\left| \begin{array}{c} p^0 \\ \swarrow \searrow \\ \bullet \\ \swarrow \searrow \\ f \\ + \\ p^0 \\ \swarrow \searrow \\ \bullet \\ \swarrow \searrow \\ f \end{array} \right|^2 \neq \left| \begin{array}{c} \bar{p}^0 \\ \swarrow \searrow \\ \bullet \\ \swarrow \searrow \\ \bar{f} \\ + \\ \bar{p}^0 \\ \swarrow \searrow \\ \bullet \\ \swarrow \searrow \\ \bar{f} \end{array} \right|^2$$



- SM with only a source: the phase in CKM
- in charm sector, it's predicted at $\sim \mathcal{O}(10^{-3})$
- $\sim 1\%$ exp. sensitivity to observe CPV \rightarrow NP



Status of D^0 - \bar{D}^0 mixing and CP violation [mainly ref. charm physics at HFLAV]

 D^0 - \bar{D}^0 mixing

indirect CP violation

CP violation in $D^0 \rightarrow KK/\pi\pi$ 

- $\gg 11.5\sigma$ to exclude no mixing $(x,y)=(0,0)$ with CPV-allowed
- No hints for indirect CPV \Leftarrow no direct CPV $(|q/p|,\phi)=(1,0)$ at C.L=40%
- No clear evidence of direct CPV \Leftarrow no CPV at C.L=9.3%

D^0 - \bar{D}^0 mixing observation in more channels, and CPV searches are two of most important physical goals for Charm WG at our Belle II experiment.






Improved D^0 proper time resolution at Belle II

- ▶ Time-dependent amplitude of $D^0 \rightarrow f$ (here $t[\tau_{D^0}]$ and $\lambda_f = \frac{q}{p} \frac{A_f}{\bar{A}_f}$):

$$\Gamma(D^0(t) \rightarrow f) \propto |\mathcal{A}_f|^2 e^{-t} \left(\frac{1+|\lambda_f|^2}{2} \cosh(yt) - \text{Re}(\lambda_f) \sinh(yt) \frac{1-|\lambda_f|^2}{2} \cos(xt) + \text{Im}(\lambda_f) \sin(xt) \right) \otimes_t \text{Res}(t)$$

- ▶ Time resolution $\text{Res}(t)$ is essential in t-dept. measurements of D^0 - \bar{D}^0 mixing and CPV
- ▶ Determine D^0 proper time: $t = \frac{\ell_{\text{dec}}}{c\beta\gamma} = \frac{m_D}{cp} \vec{d} \cdot \frac{\vec{p}}{p}$ and its uncertainty σ_t



- ▶ Based on MC study, time resolution = 140 fs: 2x better than BaBar (270 fs)
- ▶ Time error σ_t : factor 3 improvement; and $\text{RMS}(\sigma_t)$: reduced by a factor 2.
 - $\text{Res} = \text{Gauss}(\mu, k\sigma_t)$, so reduced $\text{RMS}(\sigma_t)$ (higher weight in the fit) results in an increased statistics



Hadronic wrong-sign decays $D^0 \rightarrow K^+ \pi^-$


- Wrong-sign (WS) decay rates with D^0 - \bar{D}^0 mixing or CPV-allowed:

$$\frac{N(D^0 \rightarrow f)}{dt} = e^{-\Gamma t} \left[R_D + \left| \frac{q}{p} \right| \sqrt{R_D} (y' \cos \phi - x' \sin \phi) (\Gamma t) + \left| \frac{q}{p} \right|^2 \frac{(x'^2 + y'^2)}{4} (\Gamma t)^2 \right]$$

$$\frac{N(D^0 \rightarrow f)}{dt} = e^{-\Gamma t} \left[R_D + \sqrt{R_D} y' (\Gamma t) + \frac{(x'^2 + y'^2)}{4} (\Gamma t)^2 \right] \quad (\text{no CPV})$$

where $x' = x \cos \delta + y \sin \delta$, $y' = y \cos \delta - x \sin \delta$ with strong phase difference δ

- Belle: first observation (5.1σ) of D^0 - \bar{D}^0 mixing in e^+e^- collisions [PRL 112, 111801 (2014)]
- Belle II sensitivity estimation based on ToyMC
 - Smear decay time with Gauss ($\sigma = 140$ fs) for 1000 experiments
 - Obtain sensitivities by RMS of residuals distribution of parameters



Parameter		Belle	Belle II		
		976 /fb	5 /ab	20 /ab	50 /ab
no	$\sigma(x'^2)(10^{-5})$	22	7.5	3.7	2.3
CPV	$\sigma(y')(\%)$	0.34	0.11	0.056	0.035
CPV-allowed	$\sigma(x')(\%)$		0.37	0.23	0.15
	$\sigma(y')(\%)$		0.26	0.17	0.10
	$\sigma(q/p)$		0.197	0.089	0.051
	$\sigma(\phi)(^\circ)$		15.5	9.2	5.7

one order of magnitude better than Belle measurement



Time-dependent Dalitz analyses in D three-body decays

- ▶ Time-dependent Dalitz plot (TDDP) provides an essential tool in studying D^0 - \bar{D}^0 mixing.
- ▶ Only method: sensitive to **linear order in both mixing parameters**, especially self-conjugated decays like $K_S^0 hh$ (not rotated by an unknown δ)





- ▶ TDDP fit on $D^0 \rightarrow K^+ \pi^- \pi^0$ WS decays to extract mixing par. ($x''/r_0, y''/r_0$)

$$|A_{\bar{f}}|^2 = \left[|A_{\bar{f}}^{DCS}|^2 e^{-\Gamma t} + \frac{(x^2 + y^2)}{4r_0^2} |A_{\bar{f}}^{CF}|^2 (\Gamma t)^2 e^{-\Gamma t} + \left(\frac{y''}{r_0} \text{Re}[A_{\bar{f}}^{DCS} \bar{A}_{\bar{f}}^{CF}] + \frac{x''}{r_0} \text{Im}[A_{\bar{f}}^{DCS} \bar{A}_{\bar{f}}^{CF}] \right) (\Gamma t) e^{-\Gamma t} \right] \otimes_t \text{Res}(t)$$


$$x'' = x \cos \delta_{K\rho} + y \sin \delta_{K\rho}, \quad y'' = y \cos \delta_{K\rho} - x \sin \delta_{K\rho}, \quad r_0 = |A^{CF}|/|A^{DCS}|$$

- ▶ BaBar: the evidence (3.2σ) with 384 fb^{-1} : $\sigma(x'', y'') = \left(\begin{smallmatrix} +0.57 & +0.55 \\ -0.68 & -0.64 \end{smallmatrix} \right) \%$ [PRL 103, 211801 (2009)]
- ▶ ToyMC: smear lifetime with Gauss ($\sigma=140 \text{ fs}$); without considering bkg effects.
- ▶ Sensitivity estimation: **one order of magnitude improvement than BaBar**

detailed info. see [Chin. Phys. C, 41: 023001 (2017)]



$$\sigma_x = 0.060\%$$



$$\sigma_y = 0.049\%$$

- ▶ More t-dept. measurements, like $D^0 \rightarrow K^+ K^-, \pi^+ \pi^-, D^0 \rightarrow K_S^0 \pi^+ \pi^-$ etc. in backup.
(y_{CP}, A_{Γ}) ($x, y, |\bar{q}/p|, \phi$)




Time-integrated CP asymmetry in D decays

- Time-integrated CP asymmetries are measured based on partial decay rates:

$$A_{CP}^f = \frac{\Gamma(D \rightarrow f) - \Gamma(\bar{D} \rightarrow \bar{f})}{\Gamma(D \rightarrow f) + \Gamma(\bar{D} \rightarrow \bar{f})} = a_{dir}^f + a_{ind}^f \quad \text{e.g.: in } D^0 \rightarrow K_S^0 h^+, \text{ measured asym.: } A_{raw} = A_{CP} + A_{FB} + A_{\epsilon}^{h^+} + A_{CP}^{K^0}$$

- Several measurements are performed at Belle

Channel	$\mathcal{L}(/fb)$	Current measurement		References	Belle II	LHCb
		value(%)	References		50 ab ⁻¹ (%)	50 fb ⁻¹ (%)
$D^0 \rightarrow \pi^+ \pi^-$	976	$+0.55 \pm 0.36 \pm 0.09$	PoS ICHEP2012 (2013) 353	± 0.05	± 0.03	
$D^0 \rightarrow K^+ K^-$	976	$-0.32 \pm 0.21 \pm 0.09$	PoS ICHEP2012 (2013) 353	± 0.03	± 0.03	
$D^0 \rightarrow \pi^0 \pi^0$	966	$-0.03 \pm 0.64 \pm 0.10$	PRL 112, 211601 (2014)	± 0.09		
$D^0 \rightarrow K_S^0 K_S^0$	921	$-0.02 \pm 1.53 \pm 0.17$	PRL 119, 171801 (2017)	± 0.20		
$D^0 \rightarrow K_S^0 \pi^0$	966	$-0.21 \pm 0.16 \pm 0.07$	PRL 112, 211601 (2014)	± 0.03		
$D^0 \rightarrow K_S^0 \eta$	791	$+0.54 \pm 0.51 \pm 0.16$	PRL 106, 211801 (2011)	± 0.07		
$D^0 \rightarrow K_S^0 \eta'$	791	$+0.98 \pm 0.67 \pm 0.14$	PRL 106, 211801 (2011)	± 0.09		
$D^0 \rightarrow \pi^+ \pi^- \pi^0$	532	$+0.43 \pm 0.41 \pm 1.23$	PLB 662, 102 (2008)	± 0.13		
$D^0 \rightarrow K^+ \pi^- \pi^0$	281	-0.60 ± 5.30	PRL 95, 231801 (2005)	± 0.40		
$D^0 \rightarrow K^+ \pi^- \pi^+ \pi^-$	281	$+1.8 \pm 4.4$	PRL 95, 231801 (2005)	± 0.33		
Belle Preliminary						
$D^+ \rightarrow \pi^0 \pi^+$	921	$+2.31 \pm 1.24 \pm 0.23$		± 0.40		
$D^+ \rightarrow \phi \pi^+$	955	$+0.51 \pm 0.28 \pm 0.05$	PRL 108, 071801 (2012)	± 0.04		
$D^+ \rightarrow \eta \pi^+$	791	$+1.74 \pm 1.13 \pm 0.19$	PRL 107, 221801 (2011)	± 0.14	± 0.01	
$D^+ \rightarrow \eta' \pi^+$	791	$-0.12 \pm 1.12 \pm 0.17$	PRL 107, 221801 (2011)	± 0.14		
$D^+ \rightarrow \pi^0 \pi^+ \pi^+$	977	$-0.363 \pm 0.094 \pm 0.067(3.2\sigma)$	PRL 109, 021601 (2012)	± 0.03	± 0.03	
$D^+ \rightarrow K_S^0 K^+$	977	$-0.25 \pm 0.28 \pm 0.14$	JHEP 02 (2013) 098	± 0.05		
$D_s^+ \rightarrow K_S^0 \pi^+$	673	$+5.45 \pm 2.50 \pm 0.33$	PRL 104, 181602 (2010)	± 0.29	± 0.03	
$D_s^+ \rightarrow K_S^0 K^+$	673	$+0.12 \pm 0.36 \pm 0.22$	PRL 104, 181602 (2010)	± 0.05		



- Belle II: precision of $\mathcal{O}(0.01\%)$ (down to SM level).

$$\sigma_{Belle II} = \sqrt{(\sigma_{stat}^2 + \sigma_{syst}^2) \cdot (\mathcal{L}_{Belle}/50 \text{ ab}^{-1}) + \sigma_{fired}^2}$$

- With respect to LHCb, Belle II has advantages of excellent γ and π^0 reconstruction.
- There are some other important methods at Belle II, like T-odd asymmetry measurement with neutral particle in final states (see backup)



Outline


- 1 Belle II at SuperKEKB
- 2 D^0 - \bar{D}^0 mixing and CP violation
 - Formalism and experiment status
 - Time-dependent measurements
 - Time-integrated CP asymmetry
- 3 Rare or leptonic charm decays
 - Rare radiative decays with one photon
 - Rare radiative decays with two photons
 - (Semi-) Leptonic charm decays
- 4 ROE method of D^0 flavor tag
- 5 Summary



Rare radiative decays with one photon

Rare radiative decays of $D^0 \rightarrow V\gamma$

- ▶ Radiative decays of $D^0 \rightarrow V\gamma$: dominated by long-range contribution ($\sim 10^{-5}$, whereas short-range at 10^{-8} level); direct CPV can be enhanced to exceed 1% in NP. [PRL 109,17801(2012)]
- ▶ BR measurement to test SM; observing $A_{CP}^{V\gamma} > 3\%$ is a signal of NP. [PRL 109,17801(2012)]
- ▶ Belle II sensitivity estimation for A_{CP} based on MC study:
 - similar performance of π^0 veto and resolution of signal or bkg
 - similar signal-to-bkg btw Belle and Belle II \Rightarrow scaling luminosity



- ▶ Belle II: statistical $\sigma_{A_{CP}}$ at 1 – 2% level with 50 ab^{-1} .



radiative decays	Belle A_{CP} results ^[1] 976 fb ⁻¹	Belle II uncertainty		
		5 ab ⁻¹	15 ab ⁻¹	50 ab ⁻¹
$D^0 \rightarrow \rho^0\gamma$	$+0.056 \pm 0.152 \pm 0.006$	± 0.07	± 0.04	± 0.02
$D^0 \rightarrow \phi\gamma$	$-0.094 \pm 0.066 \pm 0.001$	± 0.03	± 0.02	± 0.01
$D^0 \rightarrow \bar{K}^{*0}\gamma$	$-0.003 \pm 0.020 \pm 0.000$	± 0.01	± 0.005	± 0.003

[1] T. Nanut et al. (Belle Collaboration), Phys. Rev. Lett. **118**, 051801 (2017)





Rare radiative decay of $D^0 \rightarrow \gamma\gamma$

- FCNC process is forbidden at tree level in SM; but NP modes allow it and lead to significant enhancement on BR.
- $D^0 \rightarrow \gamma\gamma$: a sensitive probe for new physics.
- Belle: most restrictive limit (832 fb^{-1}): $BR < 8.5 \times 10^{-7}$ approaching SM prediction (10^{-8})
- This FCNC decay will be probed further at **Belle II on 50 ab^{-1} : $\sim 10^{-7} - 10^{-8}$**
- With respect to LHCb, there are still some rooms for semileptonic decays (neutrinos in final states) at Belle II.



prospects for D (semi-) Leptonic decays

- ▶ (Semi-)Leptonic charm decays involve both well-understood weak interactions physics and non-perturbative strong-interaction effects \Rightarrow test lattice QCD or measure $|V_{cd}|$ and $|V_{cs}|$.
 - leptonic decays are used to extract $|V_{cd}|f_D$ or $|V_{cs}|f_{D_s}$
 - semileptonic decays are used to extract $|V_{cd}|f_+^\pi(q^2=0)$ or $|V_{cs}|f_+^K(q^2=0)$
- Study $E_{miss}(\nu)$ in $c\bar{c} \rightarrow D_{tag}^+ D^{*-} X_{frag}^0$, where $D^{*-} \rightarrow (D_{sig} \rightarrow (h)\ell\nu)\pi$ ($\ell=e/\mu; h=K/\pi$) with $P_{miss} = P_{e^+} + P_{e^-} - P_{tag} - P_{frag} - P_\ell(-P_h)$, define U_{miss} (better resol.) or M_{miss}



- Leptonic decays yields estimation at Belle II with $50 ab^{-1}$

Mode	$D_s^- \rightarrow \mu^- \bar{\nu}$	$D^0 \rightarrow \nu \bar{\nu}$	$D^- \rightarrow \mu^- \bar{\nu}$	$D \rightarrow \pi \ell^+ \nu$
Belle	$0.91 ab^{-1}$	$0.92 ab^{-1}$		$0.28 ab^{-1}$
	492 ± 26	$(695 \pm 1) \times 10^3$		126 ± 12
Belle II	27000	38×10^6	1250	7.0×10^5
Sensitivity	improve $ V_{cs} _{best}$	7 \times better	$< 2\% V_{cd} $	comparable to BESIII






Outline

- 1 Belle II at SuperKEKB
- 2 D^0 - \bar{D}^0 mixing and CP violation
 - Formalism and experiment status
 - Time-dependent measurements
 - Time-integrated CP asymmetry
- 3 Rare or leptonic charm decays
 - Rare radiative decays with one photon
 - Rare radiative decays with two photons
 - (Semi-) Leptonic charm decays
- 4 ROE method of D^0 flavor tag
- 5 Summary



ROE: a new D^0 flavor tagging method

- To measure CPV, the flavor of D^0 is determined effectively.
- At B-factories, the charge of π_s from $D^{*+} \rightarrow D^0 \pi_s^+$ is used to tag the flavor of D^0 ; but D^0 mesons from B decays are excluded.
 \Rightarrow only D^0 from $D^{*\pm}$ in $c\bar{c}$ events (25%) were used.
- **ROE** method: select events with only one K^\pm in the **Rest Of Event**;
- the charge of this K^\pm in ROE to determine the flavor of D^0 .



$$\frac{N_{tag}^0}{N_{tag}^*} = \frac{\epsilon_{tag}^0}{\epsilon_{tag}^*} \cdot \frac{N_{gen}^0 + (1 - \epsilon_{tag}^*) \cdot N_{gen}^*}{N_{gen}^*} \sim 1$$

here ϵ_{tag}^* (ϵ_{tag}^0): tagging efficiency of D^* (ROE) method with 80% ($\leq 20\%$).

N_{gen}^* (N_{gen}^0): number of D^0 produced by a D^* (other $c\bar{c}$ event) with $N_{gen}^0 : N_{gen}^* \simeq 3 : 1$

A reduction of $\sim 15\%$ of $\sigma(stat)$ on A_{CP}

An additional D^0 sample from ROE for mixing and CPV measurements.



Outline

- 1 Belle II at SuperKEKB
- 2 D^0 - \bar{D}^0 mixing and CP violation
 - Formalism and experiment status
 - Time-dependent measurements
 - Time-integrated CP asymmetry
- 3 Rare or leptonic charm decays
 - Rare radiative decays with one photon
 - Rare radiative decays with two photons
 - (Semi-) Leptonic charm decays
- 4 ROE method of D^0 flavor tag
- 5 Summary



Summary

- ▶ Belle II at SuperKEKB which aims to achieve luminosity of $8.0 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$, will collect 50 ab^{-1} of dataset, which gives us a **rich program for charm physics study**.
- ▶ Many impacts on charm physics at Belle II with the large dataset are presented, benefiting from the improved tracking efficiency and vertex reconstruction,
 - D^0 - \bar{D}^0 mixing and CP violation measurement with much more precision
 - more precise and exciting results for CP asymmetries
 - competitive in searches of several rare charm decays
- ▶ A new D^0 flavor tagging method, **ROE**, will increase an additional D^0 sample.
- ▶ Belle II will achieve more precise measurements (mostly one order of magnitude improvement) of charm observables in the next decade, improving our knowledge of charm physics and searching for new physics beyond the Standard Model.

Let's look forwards to the charming news of charm physics from Belle II.





Thank you for your attention.


谢谢!



Longke LI (李龙科)
Room A519, Main Building
Institute of High Energy Physics, CAS (IHEP)
19B, Yuquan Road, Shijingshan District
Beijing City, 100049, P. R. China







 (+86)-159-5693-4447

 lilongke_ustc

 lilongke@ihep.ac.cn



Status of D^0 - \bar{D}^0 mixing and CP violation [mainly ref. charm physics at HFLAV]

Decay Type	Final State						
WS 2-body decay	$K^+ \pi^-$	★	☆	★	★	✓	✓ $_{\delta K\pi}$
CP-eigenstates	$K^+ K^- / \pi^+ \pi^-$	☆	☆	☆ $_{ACP}^{(a)}$	✓ $_{ACP}$	✓	
WS 3-body decay	$K^+ \pi^- \pi^0$	○ $^{(b)}$	☆			✓ $_{ACP}$	○ $_{\delta}$
Self-conjugated 3-body decay	$K_S^0 \pi^+ \pi^-$	✓	✓	✓	✓ $_{ACP}$	✓	○ $_{\delta}$
	$K_S^0 K^+ K^-$	○ $^{(c)}$	✓	○			○ $_{\delta}$
	$K_S^0 \pi^0 \pi^0$					✓ $_{Dalitz}$	○ $_{YCP}$
Self-conjugated SCS 3-body decay	$\pi^+ \pi^- \pi^0$	✓ $_{ACP}$	✓ $_{ACP}^{mixing}$	✓ $_{ACP}$			○ $_{\delta}$
	$K^+ K^- \pi^0$		✓ $_{ACP}$				○ $_{\delta}$
SCS 3-body	$K_S^0 K^{\pm} \pi^{\mp}$			✓ $_{ACP}$		✓ $_{\delta}$	○ $_{\delta}$
Semileptonic decay	$K^+ \ell^- \nu_{\ell}$	✓	✓			✓	
Multi-body($n \geq 4$)	$K^+ \pi^- \pi^+ \pi^-$	✓ $_{RWS}$	✓	★			○ $_{\delta RS}$
	$\pi^+ \pi^- \pi^+ \pi^-$	○ $_{ACP}$		✓ $_{ACP}$			
	$K^+ K^- \pi^+ \pi^-$	○ $_{AT}$	✓ $_{AT}$	✓ $_{ACP}^{(d)}$		✓ $_{ACP}$	○
	$K_S^0 \pi^+ \pi^- \pi^0$	✓ $_{AT}$					
$\psi(3770) \rightarrow D^0 \bar{D}^0$ via correlations						✓ $_{\delta K\pi}$	✓ $_{YCP}$

★ for observation ($> 5\sigma$); ☆ for evidence ($> 3\sigma$); ✓ for measurement published; ○ for analysis on going. A_T stands for measuring CP asymmetry using T-odd correlations.

The related publications are linked under their corresponding signs.

(a) LHCb also give the measurement of indirect CP asymmetry in $D^0 \rightarrow h^- h^+$ decay in PRL **112**, 041801 (2014).

(b) Belle measured WS-to-RS ratio R_{WS} and A_{CP} in $D^0 \rightarrow K^{\mp} \pi^{\pm} \pi^0$ in PRL **95**, 231801 (2005).

(c) Belle measured y_{CP} in $D^0 \rightarrow K_S^0 \phi$ in PRD **80**, 052006 (2009), the amplitude analysis for mixing parameters is on going.

(d) LHCb also search for CP violation using T-odd correlations in $D^0 \rightarrow K^+ K^- \pi^+ \pi^-$ decays in JHEP **10**(2014) 005.



Belle II detector upgrade

- Vertex detector:
 - 2 layers PXD (DEPFET) + 4 layers SVD (DSSD)
 - smaller inner radius, larger outer radius
 - ⇒ better vertex resolution
 - ⇒ improved efficiency for slow pion and K_S^0
- Central drift chamber
 - smaller cells, larger outer radius
 - ⇒ improved momentum resolution and dE/dx
- Hadron Identification
 - TOP (barrel) and Aerogel RICH (forward): replace ACC+TOF
 - ⇒ less material in front of calorimeter
 - ⇒ improved hadron ID
- Electromagnetic calorimeter
 - waveform sampling technique to cope with increase background
- K_L^0 and muon detector
 - RPC's in endcaps and first two layers of barrel replaced with scintillator counters to cope with increased neutron background



D^0 Decays to CP eigenstates

- Mixing and CPV: using CP eigenstates D^0 lifetime analysis relative to non-CP eigenstates (here $h=K/\pi$):

$$y_{CP} = \frac{\tau_{K\pi}}{\langle \tau_{hh} \rangle} - 1 \quad A_{\Gamma} = \frac{\tau(\bar{D}^0 \rightarrow hh) - \tau(D^0 \rightarrow hh)}{\tau(\bar{D}^0 \rightarrow hh) + \tau(D^0 \rightarrow hh)}$$

- if CP is conserved, $y_{CP} = y$;
- if direct CPV negligible, $y_{CP} = y \cos \phi - \frac{1}{2} A_M \sin \phi$, $A_{\Gamma} = \frac{1}{2} A_M y \cos \phi - x \sin \phi$
- Belle: the first evidence (540 fb^{-1}): $y_{CP} = (+1.31 \pm 0.32 \pm 0.25)\%$ [PRL 98, 211801 (2007)]
- Belle updated result (976 fb^{-1}) [PLB 753, 412 (2016)]

$$y_{CP} = [1.11 \pm 0.22 \pm 0.09]\% (4.7\sigma)$$


$$A_{\Gamma} = [-0.03 \pm 0.20 \pm 0.07]\%$$

- Belle II sensitivity (50 ab^{-1}):

$$\sigma_{y_{CP}} \approx 0.06\%, \quad \sigma_{A_{\Gamma}} \approx 0.04\%$$

One order of magnitude improvement

- other CP-eigenstate decays: $K_S^0 \phi$, $K_S^0 \omega$, $K_S^0 a_0(980)$ etc.









D^0 three-body self-conjugated decays

D^0 three-body self-conjugated decays $D^0 \rightarrow K_S^0 \pi \pi$

Belle $K_S^0 \pi^+ \pi^-$ by time-dependent Dalitz analysis [T. Peng *et al.* PRD 89, 091103(R) (2014)]

- direct measure x and y ; search for CPV: $q/p \neq 1$
- 921 fb⁻¹ dataset: $\sim 1.23\text{M}$ signals with high purity 95.5%
- a sum of quasi-2-body amplitudes: CF/DCS/CP-eigenstate decays, like $K^{*\mp} \pi^\pm, \rho K_S^0 \dots$
- isobar model with form factor (Blatt-Weisskopf), angular dependence (Zemach tensor)



dataset	Belle	Belle II	
	976 fb ⁻¹	5 ab ⁻¹	50 ab ⁻¹
$x(\%)$	$0.56 \pm 0.19^{+0.03+0.06}_{-0.09-0.09}$	$\pm 0.08 \pm 0.11$	$\pm 0.03 \pm 0.11$
$y(\%)$	$0.30 \pm 0.15^{+0.04+0.03}_{-0.05-0.06}$	$\pm 0.06 \pm 0.05$	$\pm 0.02 \pm 0.04$
$ q/p $	$0.90^{+0.16+0.05+0.06}_{-0.15-0.04-0.05}$	$\pm 0.07 \pm 0.07$	$\pm 0.02 \pm 0.07$
$\arg(q/p)(^\circ)$	$-6 \pm 11 \pm 3^{+3}_{-4}$	$\pm 5 \pm 4$	$\pm 2 \pm 4$



T-odd asymmetry measurements in D four-body decays

- T-odd correlations provides a powerful tool to indirectly search for CP violation:

(1) a triple product of momenta; (2) assuming CPT symmetry

- Parity-odd observable $C_T = (\vec{p}_1 \times \vec{p}_2) \cdot \vec{p}_3$ and its CP-conjugated observable \bar{C}_T

$$A_T = \frac{\Gamma(C_T > 0) - \Gamma(C_T < 0)}{\Gamma(C_T > 0) + \Gamma(C_T < 0)} \quad \bar{A}_T = \frac{\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)}{\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)}$$


- T-odd asymmetry definition to veto FSI effects:

$$a_{CP}^{T\text{-odd}} = \frac{1}{2}(A_T - \bar{A}_T) \quad \text{can be nonzero if CPV}$$

- Observing a T-odd asymmetry would be a signal for processes beyond the SM.

- Status of T-odd asymmetries in charmed mesons decay-rates:

$D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$	$a_{CP}^{T\text{-odd}} = (-0.28 \pm 1.38_{-0.76}^{+0.23}) \times 10^{-3}$	Belle ^[1]
$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$	$a_{CP}^{T\text{-odd}} = (+1.7 \pm 2.7) \times 10^{-3}$	LHCb ^[2] , BaBar ^[3] , Focus ^[4]
$D^+ \rightarrow K_S^0 K^+ \pi^+ \pi^-$	$a_{CP}^{T\text{-odd}} = (-1.10 \pm 1.09) \times 10^{-2}$	BaBar ^[5] , Focus ^[4]
$D_s^+ \rightarrow K_S^0 K^+ \pi^+ \pi^-$	$a_{CP}^{T\text{-odd}} = (-1.39 \pm 0.84) \times 10^{-2}$	BaBar ^[5] , Focus ^[4]



[1] K. Prasanth et al. (Belle Collab.), Phys. Rev. D **95**, 091101(R) (2017)
 [2] R. Aaij et al. (LHCb Collab.), JHEP **10**, 5 (2014)
 [3] P. del Amo Sanchez et al. (BaBar Collab.), Phys. Rev. D **81**, 111103(R) (2010)
 [4] J.M. Link et al. (FOCUS Collab.), Phys. Lett. B **622**, 239 (2005)
 [5] J.P. Lees et al. (BaBar Collab.), Phys. Rev. D **84**, 031103(R) (2011)

Belle II could improve these results with more precision benefited from the increased dataset.



A_{CP} in charm decays into neutral kaons [ref. Fu-Sheng Yu et al., arXiv:1707.09297]

- A new CP violation effect in charm decays into neutral kaons, e.g: $D^+ \rightarrow K_S^0 \pi^0$ with evidence for CP violation: $A_{CP} = (-0.363 \pm 0.094 \pm 0.067)\%$

Indirect CPV in kaon mixing


$$A_{CP}(t) \simeq \left[\overline{A_{CP}^{K^0}}(t) + A_{CP}^{dir}(t) + A_{CP}^{int}(t) \right] / D(t)$$

$$A_{CP}^{K^0}(t) = 2e^{-t/\tau_S} \left[\mathcal{R}e(\epsilon) - e^{\Delta t_1/2} \left(\mathcal{R}e(\epsilon) \cos(\Delta mt) + \mathcal{I}m(\epsilon) \sin(\Delta mt) \right) \right] (10^{-3})$$


CPV in interference between kaon mixing and charm decays

$$A_{CP}(t) \simeq \left[A_{CP}^{K^0}(t) + A_{CP}^{dir}(t) + \overline{A_{CP}^{int}}(t) \right] / D(t)$$

$$A_{CP}^{int}(t) = e^{-t/\tau_S} \left[4r_f \cos \phi \sin \delta_f \left[-\mathcal{I}m(\epsilon) + e^{\Delta t_1/2} \left(\mathcal{I}m(\epsilon) \cos(\Delta mt) - \mathcal{R}e(\epsilon) \sin(\Delta mt) \right) \right] \right]$$



Report at Beihang Univ. at 22nd Jun. 2017





Impacts on measurements of CP asymmetries ref. [BELLE2-NOTE-PH-2017-001]

- ▶ Ratio $\frac{\sigma_A^0}{\sigma_A^*}$ of A_{CP} statistical sensitivity σ between ROE method and D^* method

$$\alpha = \frac{\sigma_A^0}{\sigma_A^*} = \sqrt{\frac{1}{3} \cdot \frac{Q^*}{Q^0} \cdot \frac{\rho_{rec}^*}{\rho_{rec}^0}}$$

- effective tagging efficiency: $Q = \epsilon_{tag}(1 - 2\omega)^2$ with mistagging ratio ω ;
- ρ_{rec} : purity of reconstructed D .

- ▶ Combining ROE and D^* method: $\sigma_A^C = \frac{\alpha}{\sqrt{1+\alpha}} \simeq 0.85 \cdot \sigma_A^*$, with assuming $\rho_{rec}^* / \rho_{rec}^0 = 1.4$



A reduction of $\sim 15\%$ of $\sigma(stat)$ on A_{CP}

- Criteria A: basic (see ref. [BELLE2-NOTE-PH-2017-001])
- Criteria B: veto rec.ed $K_S^0 \rightarrow \pi^+ \pi^-$ in ROE; proper cut on $\cos \theta_{K^+ D^0}$
- Criteria C: veto all K_S^0 / K_L^0 gen.ed in ROE; same cut on $\cos \theta_{K^+ D^0}$ as before

